

RAR γ siRNA (m): sc-36390

BACKGROUND

Retinoids are metabolites of vitamin A (retinol) that are important signaling molecules during vertebrate development and tissue differentiation. Retinoic acid receptors (RARs) and retinoid X receptors (RXRs) are nuclear transcription factors that modulate the effects of retinoids (RA) on gene expression. Most retinoid forms (including all *trans* RA, 9-*cis* RA, 4-oxo RA and 3,4 dihydro RA) activate RAR family members, whereas RXR family members are activated by 9-*cis*-RA only. RA binds RARs, inducing a change in receptor configuration that allows DNA binding and increased gene transcription from specific genes to occur. RAR family members, which include RAR α , RAR β and RAR γ , belong to the same class of nuclear transcription factors as thyroid hormone receptors, vitamin D₃ receptor and ecdysone receptor. Retinoid receptor expression is tissue specific; the skin expresses RAR γ and RXR α . The expression of RAR γ and RXR β was somewhat decreased in lung cancers. The human RAR γ gene maps to chromosome 12q13.13.

REFERENCES

- Mattei, M.G., et al. 1991. Chromosomal assignment of retinoic acid receptor (RAR) genes in the human, mouse, and rat genomes. *Genomics* 10: 1061-1069.
- Koelle, M.R., et al. 1991. The *Drosophila* EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily. *Cell* 67: 59-77.
- Rees, J. 1992. The molecular biology of retinoic acid receptors: orphan from good family seeks home. *J. Dermatol.* 126: 97-104.
- Bhat, M.K., et al. 1994. Phosphorylation enhances the target gene sequence-dependent dimerization of thyroid hormone receptor with retinoid X receptor. *Proc. Natl. Acad. Sci. USA* 91: 7927-7931.

CHROMOSOMAL LOCATION

Genetic locus: Rarg (mouse) mapping to 15 F3.

PRODUCT

RAR γ siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RAR γ shRNA Plasmid (m): sc-36390-SH and RAR γ shRNA (m) Lentiviral Particles: sc-36390-V as alternate gene silencing products.

For independent verification of RAR γ (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-36390A, sc-36390B and sc-36390C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RAR γ siRNA (m) is recommended for the inhibition of RAR γ expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

RAR γ (G-1): sc-7387 is recommended as a control antibody for monitoring of RAR γ gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RAR γ gene expression knockdown using RT-PCR Primer: RAR γ (m)-PR: sc-36390-PR (20 μ l, 456 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Tamaki, M., et al. 2019. All-*trans* retinoic acid suppresses bone morphogenetic protein 4 in mouse diabetic nephropathy through a unique retinoic acid response element. *Am. J. Physiol. Endocrinol. Metab.* 316: E418-E431.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.