

Ribophorin I siRNA (h): sc-36420

BACKGROUND

Membrane proteins of the endoplasmic reticulum (ER) may be localized by mechanisms that involve retention, retrieval or a combination of both. ER localization information has been found in cytoplasmic, transmembrane or luminal domains. Specific retrieval mechanisms have been identified for luminal ER proteins, which contain a KDEL domain, and for type I transmembrane proteins carrying a dilysine motif. Mammalian oligosaccharyltransferase (OST) is a protein complex that is composed of four rough ER-specific, type I transmembrane proteins: Ribophorins I and II (RI and RII), OST48 and DAD1 (also designated defender against apoptotic death). The ribophorins are integral membrane glycoproteins that localize exclusively to the rough ER. There is affinity between the cytoplasmically located N-terminal region of DAD1 and the short cytoplasmic tail of OST48 to place DAD1 firmly into the OST complex. The OST complex affects the cotranslational N-glycosylation of newly synthesized polypeptides.

REFERENCES

1. Silberstein, S., et al. 1992. The 48 kDa subunit of the mammalian oligosaccharyltransferase complex is homologous to the essential yeast protein WBP1. *J. Biol. Chem.* 267: 23658-23663.
2. Fu, J., et al. 1997. Interactions among subunits of the oligosaccharyltransferase complex. *J. Biol. Chem.* 272: 29687-29692.
3. Kelleher, D.J., et al. 1997. DAD1, the defender against apoptotic cell death, is a subunit of the mammalian oligosaccharyltransferase. *Proc. Natl. Acad. Sci. USA* 94: 4994-4999.
4. Sanjay, A., et al. 1998. DAD1 is required for the function and the structural integrity of the oligosaccharyltransferase complex. *J. Biol. Chem.* 273: 26094-26099.
5. Fu, J., et al. 2000. Localization of Ribophorin II to the endoplasmic reticulum involves both its transmembrane and cytoplasmic domains. *Eur. J. Cell Biol.* 79: 219-228.
6. Fu, J. and Kreibich, G. 2000. Retention of subunits of the oligosaccharyltransferase complex in the endoplasmic reticulum. *J. Biol. Chem.* 275: 3984-3990.

CHROMOSOMAL LOCATION

Genetic locus: RPN1 (human) mapping to 3q21.3.

PRODUCT

Ribophorin I siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribophorin I shRNA Plasmid (h): sc-36420-SH and Ribophorin I shRNA (h) Lentiviral Particles: sc-36420-V as alternate gene silencing products.

For independent verification of Ribophorin I (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-36420A, sc-36420B and sc-36420C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ribophorin I siRNA (h) is recommended for the inhibition of Ribophorin I expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Ribophorin I (E-7): sc-48367 is recommended as a control antibody for monitoring of Ribophorin I gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ribophorin I gene expression knockdown using RT-PCR Primer: Ribophorin I (h)-PR: sc-36420-PR (20 μ l, 411 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.