

Integrin α 4 (A-7): sc-365209

BACKGROUND

Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. The 16α and 8β subunits heterodimerize to produce more than 20 different receptors. Most Integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, collagen and Vitronectin. Certain Integrins can also bind to soluble ligands such as Fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster Integrins by binding to adjacent Integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of Integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, Integrins function as signaling receptors. Signals transduced by Integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis.

CHROMOSOMAL LOCATION

Genetic locus: ITGA4 (human) mapping to 2q31.3.

SOURCE

Integrin α 4 (A-7) is a mouse monoclonal antibody raised against amino acids 796-1005 mapping near the C-terminus of Integrin α 4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Integrin α 4 (A-7) is available conjugated to agarose (sc-365209 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-365209 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-365209 PE), fluorescein (sc-365209 FITC), Alexa Fluor[®] 488 (sc-365209 AF488), Alexa Fluor[®] 546 (sc-365209 AF546), Alexa Fluor[®] 594 (sc-365209 AF594) or Alexa Fluor[®] 647 (sc-365209 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-365209 AF680) or Alexa Fluor[®] 790 (sc-365209 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Integrin α 4 (A-7) is recommended for detection of Integrin α 4 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Integrin α 4 siRNA (h): sc-35685, Integrin α 4 shRNA Plasmid (h): sc-35685-SH, and Integrin α 4 shRNA (h) Lentiviral Particles: sc-35685-V.


Molecular Weight of Integrin α 4: 150 kDa.

Positive Controls: MOLT-4 cell lysate: sc-2233, Jurkat whole cell lysate: sc-2204 or HeLa whole cell lysate: sc-2200.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Integrin α 4 (A-7): sc-365209. Western blot analysis of Jurkat whole cell lysate.



Integrin α 4 (A-7): sc-365209. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lymph node tissue showing membrane and cytoplasmic staining of cells in germinal and non-germinal centers (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human spleen tissue showing membrane staining of cells in white pulp and cells in red pulp (B).

SELECT PRODUCT CITATIONS

- Liu, C.C., et al. 2013. The membrane proximal KXGFFKR motif of α -Integrin mediates chemoresistance. *Mol. Cell. Biol.* 33: 4334-4345.
- Bailón, E., et al. 2014. Overexpression of progelatinase B/proMMP-9 affects migration regulatory pathways and impairs chronic lymphocytic leukemia cell homing to bone marrow and spleen. *J. Leukoc. Biol.* 96: 185-199.
- Cho, H.R., et al. 2018. Radiogenomics profiling for glioblastoma-related immune cells reveals CD49d expression correlation with MRI parameters and prognosis. *Sci. Rep.* 8: 16022.
- Krajnak, S., et al. 2022. Role of integrins in the metastatic spread of high-grade serous ovarian cancer. *Arch. Gynecol. Obstet.* 305: 1291-1298.
- Ding, Y., et al. 2022. Candesartan reduces neuronal apoptosis caused by ischemic stroke via regulating the FFAR1/ITGA4 pathway. *Mediators Inflamm.* 2022: 2356507.
- Pan, R., et al. 2022. RSP02 promotes progression of ovarian cancer through dual receptor-mediated FAK/Src signaling activation. *iScience* 25: 105184.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.