

C1s (D-6): sc-365273

BACKGROUND

The complement component proteins, C1, C3, C4, and C5, are potent anaphylatoxins that are released during complement activation. Binding of these proteins to their respective G protein-coupled receptors induces proinflammatory events, such as cellular degranulation, smooth muscle contraction, arachidonic acid metabolism, cytokine release, leukocyte activation and cellular chemotaxis. C1q, together with proenzymes C1r and C1s, yield C1, the first component of the classical pathway of the serum complement system. C1 consists of a calcium dependent trimolecular complex of C1r, C1s and C1q in a 2:2:1 ratio. Activated C1s is in the form of a disulfide-linked heterodimer consisting of a heavy chain and a light chain. Defects in the gene encoding for C1s can cause selective C1s deficiency, a disorder characterized by early onset of various autoimmune diseases.

CHROMOSOMAL LOCATION

Genetic locus: C1S (human) mapping to 12p13.31.

SOURCE

C1s (D-6) is a mouse monoclonal antibody raised against amino acids 140-208 mapping within an internal region of C1s of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

C1s (D-6) is available conjugated to agarose (sc-365273 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-365273 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-365273 PE), fluorescein (sc-365273 FITC), Alexa Fluor® 488 (sc-365273 AF488), Alexa Fluor® 546 (sc-365273 AF546), Alexa Fluor® 594 (sc-365273 AF594) or Alexa Fluor® 647 (sc-365273 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-365273 AF680) or Alexa Fluor® 790 (sc-365273 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

In addition, C1s (D-6) is available conjugated to biotin (sc-365273 B), 200 µg/ml, for WB, IHC(P) and ELISA.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

C1s (D-6) is recommended for detection of mature C1 esterase and C1s precursor of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for C1s siRNA (h): sc-60301, C1s shRNA Plasmid (h): sc-60301-SH and C1s shRNA (h) Lentiviral Particles: sc-60301-V.

Molecular Weight of C1s: 88 kDa.


Positive Controls: C1s (h): 293T Lysate: sc-116635 or human platelet extract: sc-363773.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



C1s (D-6): sc-365273. Western blot analysis of C1s expression in non-transfected: sc-117752 (**A**), human C1s transfected: sc-116635 (**B**) 293T whole cell lysates and human platelet extract (**C**).



C1s (D-6): sc-365273. Immunofluorescence staining of methanol-fixed HeLa cells showing cell surface localization.

SELECT PRODUCT CITATIONS

1. Feng, P., et al. 2021. Early pregnancy regulates expression of complement components in ovine liver. *Anim. Sci. J.* 92: e13660.
2. Zhang, L., et al. 2022. Complement regulation in ovine lymph nodes during early pregnancy. *Exp. Ther. Med.* 23: 166.
3. Zhang, L., et al. 2022. Effects of early pregnancy on the complement system in the ovine thymus. *Vet. Res. Commun.* 46: 137-145.
4. Han, X., et al. 2022. Selection of early pregnancy specific proteins and development a rapid immunochromatographic test strip in cows. *Theriogenology* 187: 127-134.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.