

# PNP (F-10): sc-365551

## BACKGROUND

Purine nucleoside phosphorylase (PNP), also designated inosine phosphorylase, forms a homotrimer. It belongs to the PNP/MTAP phosphorylase family of proteins. Human PNP catalyzes the reversible phosphorolysis of ribonucleosides and 2'-deoxyribonucleosides with specificity for guanine, hypoxanthine and their analogs. PNP deficiency is a rare autosomal recessive genetic disease associated with a severe defect in T-lymphocyte function and neurologic disorder in children, comprising four percent of combined immunodeficiency cases. Children with PNP deficiency are highly prone to infections, auto-immune disorders, neurological impairment and cancer.

## REFERENCES

1. Narayana, S.V., Bugg, C.E. and Ealick, S.E. 1997. Refined structure of purine nucleoside phosphorylase at 2.75 Å resolution. *Acta Crystallogr. D. Biol. Crystallogr.* 53: 131-142.
2. Fleischman, A., Hershfield, M.S., Toutain, S., Lederman, H.M., Sullivan, K.E., Fasano, M.B., Greene, J. and Winkelstein, J.A. 1998. Adenosine deaminase deficiency and purine nucleoside phosphorylase deficiency in common variable immunodeficiency. *Clin. Diagn. Lab. Immunol.* 5: 399-400.
3. Carlucci, F., Tabucchi, A., Aiuti, A., Rosi, F., Floccari, F., Pagani, R. and Marinello, E. 2003. Capillary electrophoresis in diagnosis and monitoring of adenosine deaminase deficiency. *Clin. Chem.* 49: 1830-1838.
4. Zang, Y., Wang, W.H., Wu, S.W., Ealick, S.E. and Wang, C.C. 2005. Identification of a subversive substrate of *Trichomonas vaginalis* purine nucleoside phosphorylase and the crystal structure of the enzyme-substrate complex. *J. Biol. Chem.* 280: 22318-22325.
5. Canduri, F., Fadel, V., Dias, M.V., Basso, L.A., Palma, M.S., Santos, D.S. and de Azevedo, W.F., Jr. 2005. Crystal structure of human PNP complexed with hypoxanthine and sulfate ion. *Biochem. Biophys. Res. Commun.* 326: 335-338.
6. Canduri, F., Fadel, V., Basso, L.A., Palma, M.S., Santos, D.S. and de Azevedo, W.F., Jr. 2005. New catalytic mechanism for human purine nucleoside phosphorylase. *Biochem. Biophys. Res. Commun.* 327: 646-649.

## CHROMOSOMAL LOCATION

Genetic locus: PNP (human) mapping to 14q11.2; Pnp2/Pnp (mouse) mapping to 14 C1.

## SOURCE

PNP (F-10) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 145-167 within an internal region of PNP of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>3</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365551 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## APPLICATIONS

PNP (F-10) is recommended for detection of PNP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PNP siRNA (h): sc-45991, PNP siRNA (m): sc-45992, PNP shRNA Plasmid (h): sc-45991-SH, PNP shRNA Plasmid (m): sc-45992-SH, PNP shRNA (h) Lentiviral Particles: sc-45991-V and PNP shRNA (m) Lentiviral Particles: sc-45992-V.


Molecular Weight of PNP: 32 kDa.

Positive Controls: rat liver extract: sc-2395, TF-1 cell lysate: sc-2412 or HL-60 whole cell lysate: sc-2209.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



PNP (F-10): sc-365551. Western blot analysis of PNP expression in HL-60 (A) and TF-1 (B) whole cell lysates and rat liver (C) and rat spleen (D) tissue extracts.



PNP (F-10): sc-365551. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.