

transferrin (D-9): sc-365871

BACKGROUND

Iron (Fe) is a tightly metabolically controlled mineral and growth factor present in all living cells. Iron not bound in erythrocyte hemoglobin is transported by transferrin (Tf), the iron transport protein of vertebrate serum. The transferrin protein contains two homologous domains, each of which contain an Fe-binding site. The majority of transferrin is synthesized in the liver and secreted into the blood, but it is also produced in lower amounts in testis and brain as well as in oligodendrocytes, where transferrin is an early marker of oligodendrocyte differentiation. From the blood, transferrin is internalized by erythroblasts and reticulocytes upon binding the transferrin receptor (TfR), also designated CD71, through a system of coated pits and vesicles. After Fe release, transferrin is returned to the extracellular medium, where it can be reused. Defects in the transferrin gene results in atransferrinemia, a rare autosomal recessive disorder characterized by microcytic anemia and iron loading.

CHROMOSOMAL LOCATION

Genetic locus: TF (human) mapping to 3q22.1.

SOURCE

transferrin (D-9) is a mouse monoclonal antibody raised against amino acids 326-390 of transferrin of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ lambda light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

transferrin (D-9) is available conjugated to agarose (sc-365871 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-365871 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-365871 PE), fluorescein (sc-365871 FITC), Alexa Fluor® 488 (sc-365871 AF488), Alexa Fluor® 546 (sc-365871 AF546), Alexa Fluor® 594 (sc-365871 AF594) or Alexa Fluor® 647 (sc-365871 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-365871 AF680) or Alexa Fluor® 790 (sc-365871 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

transferrin (D-9) is recommended for detection of transferrin of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for transferrin siRNA (h): sc-37176, transferrin shRNA Plasmid (h): sc-37176-SH and transferrin shRNA (h) Lentiviral Particles: sc-37176-V.


Molecular Weight of transferrin: 79 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, human kidney extract: sc-363764 or human heart extract: sc-363763.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGλ BP-HRP: sc-516132 or m-IgGλ BP-HRP (Cruz Marker): sc-516132-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGλ BP-FITC: sc-516185 or m-IgGλ BP-PE: sc-516186 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGλ BP-HRP: sc-516132 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



transferrin (D-9) Alexa Fluor® 790: sc-365871 AF790. Direct near-Infrared western blot analysis of transferrin expression in Hep G2 whole cell lysate (A) and human kidney (B), human heart (C) and human tonsil (D) tissue extracts. Blocked with UltraCruz® Blocking Reagent: sc-516214.



transferrin (D-9): sc-365871. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing membrane staining of hepatocytes and cytoplasmic and membrane staining of hepatic sinusoids.

SELECT PRODUCT CITATIONS

- Zhao, K.W., et al. 2015. Fibroblastic synoviocytes secrete plasma proteins via α_2 -Macroglobulins serving as intracellular and extracellular chaperones. *J. Cell. Biochem.* 116: 2563-2576.
- Staubach, S., et al. 2016. Differential proteomics of urinary exovesicles from classical galactosemic patients reveals subclinical kidney insufficiency. *J. Proteome Res.* 15: 1754-1761.
- Kim, E.K., et al. 2019. Proteomic analysis of primary colon cancer and synchronous solitary liver metastasis. *Cancer Genomics Proteomics* 16: 583-592.
- Zhao, L., et al. 2020. Serum transferrin predicts end-stage renal disease in type 2 diabetes mellitus patients. *Int. J. Med. Sci.* 17: 2113-2124.
- Sabbir, M.G., et al. 2020. Hypomorphic CAMKK2 in EA.hy926 endothelial cells causes abnormal transferrin trafficking, iron homeostasis and glucose metabolism. *Biochim. Biophys. Acta Mol. Cell Res.* 1867: 118763.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.