

# p-PKC $\delta$ (F-7): sc-365969

## BACKGROUND

Members of the protein kinase C (PKC) family play a key regulatory role in a variety of cellular functions, including cell growth and differentiation, gene expression, hormone secretion and membrane function. PKCs were originally identified as serine/threonine protein kinases whose activity was dependent on calcium and phospholipids. Diacylglycerols (DAG) and tumor promoting phorbol esters bind to and activate PKC. PKCs can be subdivided into at least two major classes, including conventional (c) PKC isoforms ( $\alpha$ ,  $\beta$ I,  $\beta$ II and  $\gamma$ ) and novel (n) PKC isoforms ( $\delta$ ,  $\epsilon$ ,  $\zeta$ ,  $\eta$  and  $\theta$ ). PKC isoforms can be activated through tyrosine phosphorylation and catalytically activated upon treatment with H2O2. The Tyr 155, 525, 523 and 565 residues in the catalytic domain are crucial for activation of these enzymes. The residue Ser 643 appears to be an autophosphorylation site.

## CHROMOSOMAL LOCATION

Genetic locus: PRKCD (human) mapping to 3p21.1; Prkcd (mouse) mapping to 14 B.

## SOURCE

p-PKC  $\delta$  (F-7) is a mouse monoclonal antibody epitope corresponding to a short amino acid sequence containing Thr 507 phosphorylated PKC  $\delta$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>3</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365969 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

p-PKC  $\delta$  (F-7) is recommended for detection of Thr 507 phosphorylated PKC  $\delta$  of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PKC  $\delta$  siRNA (h): sc-36253, PKC  $\delta$  siRNA (m): sc-36246, PKC  $\delta$  shRNA Plasmid (h): sc-36253-SH, PKC  $\delta$  shRNA Plasmid (m): sc-36246-SH, PKC  $\delta$  shRNA (h) Lentiviral Particles: sc-36253-V and PKC  $\delta$  shRNA (m) Lentiviral Particles: sc-36246-V.


Molecular Weight of p-PKC  $\delta$ : 78 kDa.

Positive Controls: RAW 264.7 whole cell lysate: sc-2211, HeLa-PMA cell lysate: sc-2258 or NIH/3T3 whole cell lysate: sc-2210.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Lambda Phosphatase: sc-200312A and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG $\kappa$  BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



Western blot analysis of PKC  $\delta$  phosphorylation in untreated (A, C) and lambda protein phosphatase (sc-200312A) treated (B, D) RAW 264.7 whole cell lysates. Antibodies tested include p-PKC  $\delta$  (F-7): sc-365969 (A, B) and PKC  $\delta$  (C-20): sc-937 (C, D).



p-PKC  $\delta$  (F-7): sc-365969. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lower stomach tissue showing cytoplasmic and membrane staining of glandular cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human lymph node tissue showing cytoplasmic and membrane staining of cells in non-germinal center (B).

## SELECT PRODUCT CITATIONS

- Hwang, Y., et al. 2021. Repeated exposure to microcystin-leucine-arginine potentiates excitotoxicity induced by a low dose of kainate. *Toxicology* 460: 152887.
- Cui, H., et al. 2021. The expression of diacylglycerol kinase isoforms  $\alpha$  and  $\zeta$  correlates with the progression of experimental autoimmune encephalomyelitis in rats. *Histochem. Cell Biol.* 156: 437-448.
- Hwang, Y., et al. 2022. Effect of rottlerin on astrocyte phenotype polarization after trimethyltin insult in the dentate gyrus of mice. *J. Neuroinflammation* 19: 142.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.