

TTF-1 siRNA (m): sc-36757

BACKGROUND

TTF-1 (thyroid transcription factor-1, BCH, BHC, NK-2, Nkx2.1, Nkx2A, TEBP, TTF1) is a member of the Nkx2 family of homeodomain-containing transcription factors and regulates the transcriptional activity of thyroid-specific genes. TTF-1 influences organogenesis and the maintenance of the differentiated phenotypes of various tissues including thyroid, lung and brain. TTF-1, which is present in the epithelium of the lung, regulates transcription of the surfactant proteins (SP) A, B and C and is essential for lung morphogenesis. In the thyroid, TTF-1 elevates the expression of thyroid specific markers, thyroglobulin, thyroperoxidase and thyrotropin receptors. TTF-1 interacts with SRC-1 and CBP *in vitro*.

REFERENCES

- Zannini, M., et al. 1996. Mapping and functional role of phosphorylation sites in the thyroid transcription factor-1 (TTF-1). *J. Biol. Chem.* 271: 2249-2254.
- Ohe, K., et al. 1996. Interferon- γ suppresses thyrotropin receptor promoter activity by reducing thyroid transcription factor-1 (TTF-1) binding to its recognition site. *Mol. Endocrinol.* 10: 826-836.
- Nakazato, M., et al. 1997. Transcription of the thyroid transcription factor-1 (TTF-1) gene from a newly defined start site: positive regulation by TTF-1 in the thyroid. *Biochem. Biophys. Res. Commun.* 238: 748-752.
- Oguchi, H., et al. 1998. Multiple transcripts encoded by the thyroid-specific enhancer-binding protein (T/EBP)/thyroid-specific transcription factor-1 (TTF-1) gene: evidence of autoregulation. *Endocrinology* 139: 1999-2006.
- Katoh, R., et al. 2000. Expression of thyroid transcription factor-1 (TTF-1) in human C cells and medullary thyroid carcinomas. *Hum. Pathol.* 31: 386-393.
- Nakazato, M., et al. 2000. Thyroglobulin repression of thyroid transcription factor 1 (TTF-1) gene expression is mediated by decreased DNA binding of nuclear factor I proteins which control constitutive TTF-1 expression. *Mol. Cell. Biol.* 20: 8499-8512.
- Gereben, B., et al. 2001. The human, but not rat, DIO2 gene is stimulated by thyroid transcription factor-1 (TTF-1). *Mol. Endocrinol.* 15: 112-124.

CHROMOSOMAL LOCATION

Genetic locus: Nkx2-1 (mouse) mapping to 12 C1.

PRODUCT

TTF-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TTF-1 shRNA Plasmid (m): sc-36757-SH and TTF-1 shRNA (m) Lentiviral Particles: sc-36757-V as alternate gene silencing products.

For independent verification of TTF-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-36757A, sc-36757B and sc-36757C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TTF-1 siRNA (m) is recommended for the inhibition of TTF-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TTF-1 (8G7G3/1): sc-53136 is recommended as a control antibody for monitoring of TTF-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TTF-1 gene expression knockdown using RT-PCR Primer: TTF-1 (m)-PR: sc-36757-PR (20 μ l, 489 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Jia, S., et al. 2020. Nkx2.1 downregulation is involved in brain abnormality induced by excess retinoic acid. *Acta Biochim. Biophys. Sin.* 52: 683-690.

RESEARCH USE

For research use only, not for use in diagnostic procedures.