



Sall1 (h): 293T Lysate: sc-369360

BACKGROUND

Sall1 (Sall1, sal-like 1, TBS, HSAL1) and Sall2 (Sall2, sal-like 2, HSAL2, p150 (Sal2)) are mammalian homologs of the *Drosophila* region-specific homeotic gene spalt (sal), which encodes a zinc finger-containing transcription regulator. *Drosophila* spalt (sal) is an essential genetic component required for the specification of posterior head and anterior tail as opposed to trunk. Mammalian Sall1 may mediate higher order chromatin structure and may be a component of a distinct heterochromatin-dependent silencing process. Sall1 is present in kidney, brain and liver. Sall2 is a p53-independent regulator of p21 and Bax, and can function in some cell types as a regulator of cell growth and survival. Human Sall2 is present in brain, heart, kidney or pancreas. Sall1 and Sall2 are expressed in different areas of the fetal brain that may represent distinct sets of neurons.

REFERENCES

1. Nielsen, T.O., et al. 2003. Tissue microarray validation of epidermal growth factor receptor and Sall2 in synovial sarcoma with comparison to tumors of similar histology. *Am. J. Pathol.* 163: 1449-1456.
2. Sato, A., et al. 2003. Zinc finger protein Sall2 is not essential for embryonic and kidney development. *Mol. Cell. Biol.* 23: 62-69.
3. Wabbels, B.K., et al. 2004. Clinical and molecular genetic findings in isolated sporadic Duane syndrome. *Klin. Monatsbl. Augenheilkd.* 221: 849-853.
4. Wabbels, B.K., et al. 2004. No evidence of Sall4-mutations in isolated sporadic duane retraction "syndrome" (DURS). *Am. J. Med. Genet.* 131: 216-218.
5. Borozdin, W., et al. 2004. Novel mutations in the gene Sall4 provide further evidence for acro-renal-ocular and Okihiro syndromes being allelic entities, and extend the phenotypic spectrum. *J. Med. Genet.* 41: e102.
6. Borozdin, W., et al. 2004. Sall4 deletions are a common cause of Okihiro and acro-renal-ocular syndromes and confirm haploinsufficiency as the pathogenic mechanism. *J. Med. Genet.* 41: e113.
7. Kohlhasse, J., et al. 2004. Mutations in Sall4 in malformed father and daughter postulated previously due to reflect mutagenesis by thalidomide. *Birth Defects Res. Part A Clin. Mol. Teratol.* 70: 550-551.

CHROMOSOMAL LOCATION

Genetic locus: SALL1 (human) mapping to 16q12.1.

PRODUCT

Sall1 (h): 293T Lysate represents a lysate of human Sall1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Sall1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Sall1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.