



HEN1 (h2): 293T Lysate: sc-369687

BACKGROUND

The helix-loop-helix (HLH) structures are known motifs commonly found in membrane-active and DNA-binding proteins. The helix-loop-helix proteins HEN1 and HEN2 are DNA-binding proteins that may be involved in cell-type determination in the early nervous system. Studies of expression in normal tissues have demonstrated expression of NHLH1/NSCL-1 and NHLH2/NSCL-2, the genes encoding HEN1 and HEN2, in the developing central and peripheral nervous system, specifically in developing neurons.

REFERENCES

1. Begley, C.G., Visvader, J., Green, A.R., Aplan, P.D., Metcalf, D., Kirsch, I.R. and Gough, N.M. 1991. Molecular cloning and chromosomal localization of the murine homolog of the human helix-loop-helix gene SCL. *Proc. Natl. Acad. Sci. USA* 88: 869-873.
2. Begley, C.G., Lipkowitz, S., Göbel, V., Mahon, K.A., Bertness, V., Green, A.R., Gough, N.M. and Kirsch, I.R. 1992. Molecular characterization of NSCL, a gene encoding a helix-loop-helix protein expressed in the developing nervous system. *Proc. Natl. Acad. Sci. USA* 89: 38-42.
3. Göbel, V., Lipkowitz, S., Kozak, C.A. and Kirsch, I.R. 1992. NSCL-2: a basic domain helix-loop-helix gene expressed in early neurogenesis. *Cell Growth Differ.* 3: 143-148.
4. Lipkowitz, S., Göbel, V., Varterasian, M.L., Nakahara, K., Tchorz, K. and Kirsch, I.R. 1992. A comparative structural characterization of the human NSCL-1 and NSCL-2 genes. Two basic helix-loop-helix genes expressed in the developing nervous system. *J. Biol. Chem.* 267: 21065-21071.
5. Ibrahim, H.R., Thomas, U. and Pellegrini, A. 2001. A helix-loop-helix peptide at the upper lip of the active site cleft of lysozyme confers potent antimicrobial activity with membrane permeabilization action. *J. Biol. Chem.* 276: 43767-43774.

CHROMOSOMAL LOCATION

Genetic locus: NHLH1 (human) mapping to 1q23.2.

PRODUCT

HEN1 (h2): 293T Lysate represents a lysate of human HEN1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

HEN1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive HEN1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.