



COLEC11 (h): 293T Lysate: sc-370384

BACKGROUND

COLEC11 (collectin sub-family member 11), also known as CLK1, CL-K1-I, MGC3279, CL-K1-II, CL-K1-IIa or CL-K1-IIb, is a 270 amino acid C-type lectin protein that contains a collagen-like domain and a carbohydrate recognition domain, and plays an important role in host-defense. COLEC11 binds to various sugars and LPS (lipopolysaccharides), which include fucose but does not bind to glucose, hnRNP, β -1,3-Gal-T3 or mannose. COLEC11 is ubiquitously expressed in most tissues with high expression in kidney, liver, fetal liver, small intestine, thymus, spinal cord, placenta, adrenal gland, pancreas and several cell lines. COLEC11 is a secreted protein and all alternatively spliced isoforms of COLEC11 have oligomeric structures created through disulfide bonding.

REFERENCES

1. Zelensky, A.N. and Gready, J.E. 2004. C-type lectin-like domains in *Fugu rubripes*. BMC Genomics 5: 51.
2. Keshi, H., Sakamoto, T., Kawai, T., Ohtani, K., Katoh, T., Jang, S.J., Motomura, W., Yoshizaki, T., Fukuda, M., Koyama, S., Fukuzawa, J., Fukuoh, A., Yoshida, I., Suzuki, Y. and Wakamiya, N. 2006. Identification and characterization of a novel human collectin CL-K1. Microbiol. Immunol. 50: 1001-1013.
3. Nieländer, I., Bug, S., Richter, J., Giefing, M., Martín-Subero, J.I. and Siebert, R. 2007. Combining array-based approaches for the identification of candidate tumor suppressor loci in mature lymphoid neoplasms. APMIS 115: 1107-1134.
4. Motomura, W., Yoshizaki, T., Ohtani, K., Okumura, T., Fukuda, M., Fukuzawa, J., Mori, K., Jang, S.J., Nomura, N., Yoshida, I., Suzuki, Y., Kohgo, Y. and Wakamiya, N. 2008. Immunolocalization of a novel collectin CL-K1 in murine tissues. J. Histochem. Cytochem. 56: 243-252.
5. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612502. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Kelley, J.M., Edberg, J.C. and Kimberly, R.P. 2010. Pathways: Strategies for susceptibility genes in SLE. Autoimmun. Rev. 9: 473-476.
7. Hsu, Y.H., Zillikens, M.C., Wilson, S.G., Farber, C.R., Demissie, S., Soranzo, N., Bianchi, E.N., Grundberg, E., Liang, L., Richards, J.B., Estrada, K., Zhou, Y., van Nas, A., Moffatt, M.F., Zhai, G., Hofman, A., et al. 2010. An integration of genome-wide association study and gene expression profiling to prioritize the discovery of novel susceptibility loci for osteoporosis-related traits. PLoS Genet. 6: e1000977.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: COLEC11 (human) mapping to 2p25.3.

PRODUCT

COLEC11 (h): 293T Lysate represents a lysate of human COLEC11 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

COLEC11 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive COLEC11 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.