

Prolactin siRNA (m): sc-37215

BACKGROUND

The anterior pituitary secretes a variety of hormones that are involved in cell growth, differentiation and development. Prolactin, a 226 amino acid protein, plays a role in multiple processes, including cell growth, reproduction and immune function. Full length Prolactin, as well as an alternative splice product lacking the third exon, are secreted by endothelial cells involved in angiogenesis. In addition to its role in mammary development and lactation, Prolactin is known to play a role in the development of mammary cancer, acting as both a mitogen and a differentiating agent. Prolactin has also been shown to enhance the proliferation of B cell hybridomas, leading to an overall increase in antibody production. In addition, Prolactin has been demonstrated to reverse the antiproliferative effects of the immunosuppressive cytokine TGF- β . Prolactin is also associated with a variety of autoimmune diseases, including arthritis and type 1 diabetes.

REFERENCES

- Goffin, V., et al. 1998. Prolactin: a hormone at the crossroads of neuro-immunoendocrinology. *Ann. N.Y. Acad. Sci.* 840: 498-509.
- Clapp, C., et al. 1998. Expression of Prolactin mRNA and of Prolactin-like proteins in endothelial cells: evidence for autocrine effects. *J. Endocrinol.* 158: 137-144.
- Vonderhaar, B.K. 1998. Prolactin: the forgotten hormone of human breast cancer. *Pharmacol. Ther.* 79: 169-178.
- Richards, S.M., et al. 1998. Prolactin is an antagonist of TGF β activity and promotes proliferation of murine B cell hybridomas. *Cell. Immunol.* 184: 85-91.
- Neidhart, M. 1998. Prolactin in autoimmune diseases. *Proc. Soc. Exp. Biol. Med.* 217: 408-419.
- Cohen, A.D., et al. 2000. Prolactin serum level in patients with breast cancer. *Isr. Med. Assoc. J.* 2: 287-289.
- Yazawa, T., et al. 2000. Prolactin induces apoptosis in the penultimate spermatogonial stage of the testes in Japanese red-bellied newt (*Cynops pyrrhogaster*). *Endocrinology* 141: 2027-2032.
- Pearce, S., et al. 2003. Prolactin, Prolactin receptor and uncoupling proteins during fetal and neonatal development. *Proc. Nutr. Soc.* 62: 421-427.

CHROMOSOMAL LOCATION

Genetic locus: Prl (mouse) mapping to 13 A3.1.

PRODUCT

Prolactin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Prolactin shRNA Plasmid (m): sc-37215-SH and Prolactin shRNA (m) Lentiviral Particles: sc-37215-V as alternate gene silencing products.

For independent verification of Prolactin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-37215A, sc-37215B and sc-37215C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Prolactin siRNA (m) is recommended for the inhibition of Prolactin expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Prolactin (H-12): sc-271773 is recommended as a control antibody for monitoring of Prolactin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Prolactin gene expression knockdown using RT-PCR Primer: Prolactin (m)-PR: sc-37215-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.