

TNF β siRNA (h): sc-37218

BACKGROUND

Tumor necrosis factor β (TNF β), also known as lymphotoxin, is a pleiotropic cytokine. TNF α , also known as cachectin, is a smaller cytokine that binds to the same receptors, producing a vast array of effects similar to those of TNF β . TNF β and TNF α share 30% amino acid homology and have similar biological activities. TNF β is produced by activated lymphocytes, including CD4⁺ T helper cell type 1 lymphocytes, CD8⁺ lymphocytes and certain B lymphoblastoid cell lines. TNF α is produced by several different cell types, including lymphocytes, neutrophils and macrophages. TNF α and TNF β can modulate many immune and inflammatory functions, while having the ability to inhibit tumor growth. Target tumor cells must express TNF receptors 1 and 2 to be killed, with the p55 receptor mediating the cytotoxic response.

REFERENCES

1. Nedwin, G.E., et al. 1985. Human lymphotoxin and tumor necrosis factor genes: structure, homology and chromosomal localization. *Nucleic Acids Res.* 13: 6361-6373.
2. Aggarwal, B.B., et al. 1985. Human tumor necrosis factor. Production, purification, and characterization. *J. Biol. Chem.* 260: 2345-2354.
3. Vilcek, J., et al. 1991. Tumor necrosis factor. New insights into the molecular mechanisms of its multiple actions. *J. Biol. Chem.* 266: 7313-7316.
4. Tartaglia, L.A., et al. 1993. Tumor necrosis factor's cytotoxic activity is signaled by the p55 TNF receptor. *Cell* 73: 213-216.
5. De Togni, P., et al. 1994. Abnormal development of peripheral lymphoid organs in mice deficient in lymphotoxin. *Science* 264: 703-707.
6. Qin, Z., et al. 1995. Tumor growth inhibition mediated by lymphotoxin: evidence of B lymphocyte involvement in the antitumor response. *Cancer Res.* 55: 4747-4751.
7. Sarin, A., et al. 1995. Cytotoxic effect of TNF and lymphotoxin on T lymphoblasts. *J. Immunol.* 155: 3716-3718.
8. Sriskandan, S., et al. 1996. Lymphotoxin- α (TNF β) during sepsis. *Cytokine* 8: 933-937.
9. Pandey, J.P., et al. 1999. TNF α and TNF β gene polymorphisms in systemic sclerosis. *Hum. Immunol.* 60: 1128-1130.

CHROMOSOMAL LOCATION

Genetic locus: LTA (human) mapping to 6p21.33.

PRODUCT

TNF β siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TNF β shRNA Plasmid (h): sc-37218-SH and TNF β shRNA (h) Lentiviral Particles: sc-37218-V as alternate gene silencing products.

For independent verification of TNF β (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-37218A, sc-37218B and sc-37218C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TNF β siRNA (h) is recommended for the inhibition of TNF β expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TNF β (E-6): sc-28345 is recommended as a control antibody for monitoring of TNF β gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TNF β gene expression knockdown using RT-PCR Primer: TNF β (h)-PR: sc-37218-PR (20 μ l, 458 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.