

CUL-5 (F-6): sc-373822



The Power to Question

BACKGROUND

Cullin proteins comprise a distinct family of mediators that participate in the selective targeting of proteins for ubiquitin (Ub)-mediated proteolysis. CUL-1, which is the mammalian homolog of Cdc53 from yeast, is an integral component of the E3 ubiquitin ligase complex designated SCF. The SCF (Skp1/CUL-1/F-box protein complex) consists of Skp1 associating with both CUL-1 and an F-box protein, such as Skp2, which determines the substrate specificity of the complex. CUL-1 mediated ubiquitination results in the degradation of cell cycle proteins cyclin D, p21 and cyclin E. Another Cullin, CUL-3 facilitates the degradation of cyclin E independent of SCF activity, while CUL-2 associates with the tumor suppressing protein VHL and Elongin B to form VBC complexes, which structurally resemble the SCF ligase. Proteolysis also occurs by way of CUL-4 associating with Nedd-8, a ubiquitin-like protein, where it too functions as an active component of a multifunctional E3 complex. CUL-5, or vasopressin-activated, calcium-mobilizing protein (VACM-1), is also included in the cullin family as it shares substantial sequence homology with CUL-1.

CHROMOSOMAL LOCATION

Genetic locus: CUL5 (human) mapping to 11q22.3; Cul5 (mouse) mapping to 9 A5.3.

SOURCE

CUL-5 (F-6) is a mouse monoclonal antibody raised against amino acids 1-300 mapping at the N-terminus of CUL-5 (cullin-5) of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CUL-5 (F-6) is available conjugated to agarose (sc-373822 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-373822 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-373822 PE), fluorescein (sc-373822 FITC), Alexa Fluor® 488 (sc-373822 AF488), Alexa Fluor® 546 (sc-373822 AF546), Alexa Fluor® 594 (sc-373822 AF594) or Alexa Fluor® 647 (sc-373822 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-373822 AF680) or Alexa Fluor® 790 (sc-373822 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

CUL-5 (F-6) is recommended for detection of CUL-5 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


CUL-5 (F-6) is also recommended for detection of CUL-5 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for CUL-5 siRNA (h): sc-37574, CUL-5 siRNA (m): sc-37575, CUL-5 shRNA Plasmid (h): sc-37574-SH, CUL-5 shRNA Plasmid (m): sc-37575-SH, CUL-5 shRNA (h) Lentiviral Particles: sc-37574-V and CUL-5 shRNA (m) Lentiviral Particles: sc-37575-V.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



CUL-5 (F-6): sc-373822. Western blot analysis of CUL-5 expression in HeLa (A), SKBR-3 (B), T-47D (C) and MOLT-4 (D) whole cell lysates.



CUL-5 (F-6): sc-373822. Near-infrared western blot analysis of CUL-5 expression in F9 (A) and MOLT-4 (B) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgGx BP-CFL 790: sc-516181.

SELECT PRODUCT CITATIONS

1. Uematsu, K., et al. 2016. ASB7 regulates spindle dynamics and genome integrity by targeting DDA3 for proteasomal degradation. *J. Cell Biol.* 215: 95-106.
2. Zhou, H., et al. 2018. High-affinity peptidomimetic inhibitors of the DCN1-UBC12 protein-protein interaction. *J. Med. Chem.* 61: 1934-1950.
3. Zhao, G., et al. 2019. Cullin5 deficiency promotes small-cell lung cancer metastasis by stabilizing Integrin β1. *J. Clin. Invest.* 129: 972-987.
4. Yu, Q., et al. 2020. Gossypol inhibits cullin neddylation by targeting SAG-CUL5 and RBX1-CUL1 complexes. *Neoplasia* 22: 179-191.
5. Mayor-Ruiz, C., et al. 2020. Rational discovery of molecular glue degraders via scalable chemical profiling. *Nat. Chem. Biol.* 16: 1199-1207.
6. Zhao, X., et al. 2021. NPRL2 reduces the niraparib sensitivity of castration-resistant prostate cancer via interacting with UBE2M and enhancing neddylation. *Exp. Cell Res.* 403: 112614.
7. Zhang, Z., et al. 2021. HIV-2 Vif and foamy virus Bet antagonize APOBEC3B by different mechanisms. *Virology* 554: 17-27.
8. Jia, Y., et al. 2021. Th1 cytokine interferon gamma improves response in HER2 breast cancer by modulating the ubiquitin proteasomal pathway. *Mol Ther.* 29: 1541-1556.
9. Gong, L., et al. 2021. CUL5-ASB6 complex promotes p62/SQSTM1 ubiquitination and degradation to regulate cell proliferation and autophagy. *Front. Cell Dev. Biol.* 9: 684885.
10. Xu, T., et al. 2022. A small molecule inhibitor of the UBE2F-CRL5 axis induces apoptosis and radiosensitization in lung cancer. *Signal Transduct. Target. Ther.* 7: 354.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA