

SIRP- α / β 1 (C-8): sc-373896

BACKGROUND

SIRPs (signal-regulatory proteins) are a family of transmembrane glycoproteins that were identified by their association with the Src homology 2 domain-containing protein-tyrosine phosphatase SHP-2 in response to Insulin. The SIRP family negatively regulates the PI 3-K pathway, which may diminish EGFR-mediated motility and survival phenotypes that contribute to transformation of certain cell types. SIRP- α is a transmembrane protein which contains an extracellular portion with three immunoglobulin-like structures and a cytoplasmic region with four potential tyrosine phosphorylation sites. SIRP- α is a substrate for activated receptor tyrosine kinases. In its tyrosine phosphorylated form, SIRP- α binds to SH-PTP2 through SH2 interactions and acts as an SH-PTP2 substrate. SIRP- α has been shown to have negative regulatory effects on cellular responses induced by growth factors, oncogenes and insulin. SIRP- β 1 shares extensive sequence homology with SIRP- α in its extracellular portion but lacks the cytoplasmic portion. SIRP- γ , originally designated SIRP- β 2 (SIRP-B2, CD172g) has unique characteristics from both the α and β versions. SIRP- γ is expressed on the majority of T cells and a proportion of B cells. CD47 associates with SIRP- γ , and this interaction signals unidirectionally only.

REFERENCES

1. Yamauchi, K., et al. 1995. Identification of the major SHPTP2-binding protein that is tyrosine-phosphorylated in response to Insulin. *J. Biol. Chem.* 270: 17716-17722.
2. Fujioka, Y., et al. 1996. A novel membrane glycoprotein, SHPS-1, that binds the SH2-domain-containing tyrosine phosphatase SHP-2 in response to mitogens and cell adhesion. *Mol. Cell. Biol.* 16: 6887-6899.

CHROMOSOMAL LOCATION

Genetic locus: SIRPA/SIRPB1 (human) mapping to 20p13.

SOURCE

SIRP- α / β 1 (C-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 22-53 near the N-terminus of SIRP- α of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-373896 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

SIRP- α / β 1 (C-8) is recommended for detection of SIRP- α and SIRP- β 1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SIRP- α / β 1/ γ siRNA (h): sc-36492, SIRP- α / β 1/ γ shRNA Plasmid (h): sc-36492-SH and SIRP- α / β 1/ γ shRNA (h) Lentiviral Particles: sc-36492-V.

Molecular Weight of unglycosylated SIRP- α : 65 kDa.

Molecular Weight of glycosylated SIRP- α : 100-150 kDa.


Molecular Weight of SIRP- β : 55 kDa.

Positive Controls: THP-1 cell lysate: sc-2238, AML-193 whole cell lysate: sc-364182 or SIRP- α (h): 293T Lysate: sc-159295.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



SIRP- α / β 1 (C-8): sc-373896. Western blot analysis of SIRP- α expression in non-transfected: sc-117752 (A) and human SIRP- α transfected: sc-159295 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

1. Bener, G., et al. 2016. Silencing of CD47 and SIRP α by polypurine reverse Hoogsteen hairpins to promote MCF-7 breast cancer cells death by PMA-differentiated THP-1 cells. *BMC Immunol.* 17: 32.



See **SIRP- α / β (A-1): sc-17803** for SIRP- α / β antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.