


SGTA (G-11): sc-373978



The Power to Question

BACKGROUND

SGTA (small glutamine-rich tetratricopeptide repeat (TPR)-containing protein A or α), also known as SGT, hSGT or UBP (Vpu-binding protein), is a ubiquitously expressed protein that contains three TPR protein-protein interaction repeats. SGTA is believed to function as a component of the androgen receptor (AR)-chaperone-co-chaperone complex, acting as a co-chaperone involved in androgen signaling. More specifically, SGTA binds to the hinge region of the AR functions to retain the AR in the cytoplasm, thereby inhibiting androgen signaling. In addition, SGTA functions as a co-chaperone for HSP 90 and HSP 70, two proteins known to participate in apoptosis. On the basis of its role in apoptosis and androgen signaling, SGTA is a potential candidate for PCOS (polycystic ovary syndrome), a disorder characterized by androgen excess, obesity and menstrual disturbances. SGTA also interacts with the nonstructural parvovirus protein NS1 and the HIV-1 proteins Vpu and $G_{\alpha\gamma}$.

REFERENCES

- Kordes, E., et al. 1998. Isolation and characterization of human SGT and identification of homologues in *Saccharomyces cerevisiae* and *Caenorhabditis elegans*. *Genomics* 52: 90-94.
- Tobaben, S., et al. 2001. A trimeric protein complex functions as a synaptic chaperone machine. *Neuron* 31: 987-999.
- Angeletti, P.C., et al. 2002. Small glutamine-rich protein/viral protein U-binding protein is a novel cochaperone that affects heat shock protein 70 activity. *Cell Stress Chaperones* 7: 258-268.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603419. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Tobaben, S., et al. 2003. A brain-specific isoform of small glutamine-rich tetratricopeptide repeat-containing protein binds to Hsc70 and the cysteine string protein. *J. Biol. Chem.* 278: 38376-38383.
- Liou, S.T., et al. 2005. Small glutamine-rich tetratricopeptide repeat-containing protein is composed of three structural units with distinct functions. *Arch. Biochem. Biophys.* 435: 253-263.

CHROMOSOMAL LOCATION

Genetic locus: SGTA (human) mapping to 19p13.3; Sgta (mouse) mapping to 10 C1.

SOURCE

SGTA (G-11) is a mouse monoclonal antibody raised against amino acids 41-120 mapping near the N-terminus of SGTA of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SGTA (G-11) is recommended for detection of SGTA of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SGTA siRNA (h): sc-97627, SGTA siRNA (m): sc-153427, SGTA shRNA Plasmid (h): sc-97627-SH, SGTA shRNA Plasmid (m): sc-153427-SH, SGTA shRNA (h) Lentiviral Particles: sc-97627-V and SGTA shRNA (m) Lentiviral Particles: sc-153427-V.


Molecular Weight of SGTA: 35 kDa.

Positive Controls: HEL 92.1.7 cell lysate: sc-2270, IMR-32 cell lysate: sc-2409 or Neuro-2A whole cell lysate: sc-364185.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



SGTA (G-11): sc-373978. Western blot analysis of SGTA expression in HEL 92.1.7 (A), IMR-32 (B), Neuro-2A (C), EOC 20 (D), M1 (E) and PC-12 (F) whole cell lysates.



SGTA (G-11): sc-373978. Immunoperoxidase staining of formalin fixed, paraffin-embedded human salivary gland tissue showing cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

- Philp, L.K., et al. 2016. Small glutamine-rich tetratricopeptide repeat-containing protein α (SGTA) ablation limits offspring viability and growth in mice. *Sci. Rep.* 6: 28950.

RESEARCH USE

For research use only, not for use in diagnostic procedures.