

HPRT (F-12): sc-373979

BACKGROUND

HPRT (hypoxanthine phosphoribosyltransferase 1), also known as HGPRT or HPRT1, is a 218 amino acid cytoplasmic protein that belongs to the purine/pyrimidine phosphoribosyltransferase family. Involved in purine metabolism, HPRT functions as a purine salvage enzyme that catalyzes the conversion of hypoxanthine and guanine to their respective mononucleotides (inosine monophosphate and guanosine monophosphate, respectively). HPRT exists as a homotetramer that can bind two magnesium ions as cofactors. Defects in the gene encoding HPRT are the cause of gout and Lesch-Nyhan syndrome (LNS), both of which are characterized by a partial or complete lack of HPRT enzymatic activity. While a partial loss of HPRT enzymatic activity results in a buildup of uric acid (gout), a total loss of enzymatic activity results in hyperuricaemia, mental retardation, choreoathetosis and compulsive self-mutilation, all of which are symptoms associated with LNS. The severity of these diseases suggests an essential role for HPRT in purine metabolism.

REFERENCES

1. Stout, J.T., et al. 1985. HPRT: gene structure, expression, and mutation. *Annu. Rev. Genet.* 19: 127-148.
2. Fujimori, S., et al. 1997. An asymptomatic germline missense base substitution in the hypoxanthine phosphoribosyltransferase (HPRT) gene that reduces the amount of enzyme in humans. *Hum. Genet.* 99: 8-10.
3. Mizunuma, M., et al. 2001. A recurrent large Alu-mediated deletion in the hypoxanthine phosphoribosyltransferase (HPRT1) gene associated with Lesch-Nyhan syndrome. *Hum. Mutat.* 18: 435-443.
4. Koina, E., et al. 2005. An inactive X specific replication origin associated with a matrix attachment region in the human X linked HPRT gene. *J. Cell. Biochem.* 95: 391-402.
5. Dawson, P.A., et al. 2005. Normal HPRT coding region in a male with gout due to HPRT deficiency. *Mol. Genet. Metab.* 85: 78-80.
6. Cossu, A., et al. 2006. HPRTSardinia: a new point mutation causing HPRT deficiency without Lesch-Nyhan disease. *Biochim. Biophys. Acta* 1762: 29-33.
7. Keebaugh, A.C., et al. 2007. Gene duplication and inactivation in the HPRT gene family. *Genomics* 89: 134-142.
8. Liu, S., et al. 2008. HPRT mutations in lymphocytes from 1,3-butadiene-exposed workers in China. *Environ. Health Perspect.* 116: 203-208.

CHROMOSOMAL LOCATION

Genetic locus: HPRT1 (human) mapping to Xq26.2; Hprt (mouse) mapping to X A5.

SOURCE

HPRT (F-12) is a mouse monoclonal antibody raised against amino acids 1-218 representing full length HPRT of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

HPRT (F-12) is recommended for detection of HPRT of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HPRT siRNA (h): sc-40679, HPRT siRNA (m): sc-40680, HPRT shRNA Plasmid (h): sc-40679-SH, HPRT shRNA Plasmid (m): sc-40680-SH, HPRT shRNA (h) Lentiviral Particles: sc-40679-V and HPRT shRNA (m) Lentiviral Particles: sc-40680-V.


Molecular Weight of HPRT: 23 kDa.

Positive Controls: MIA PaCa-2 cell lysate: sc-2285, Hep G2 cell lysate: sc-2227 or A549 cell lysate: sc-2413.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



HPRT (F-12): sc-373979. Western blot analysis of HPRT expression in Hep G2 (A), KNRK (B), HeLa (C), A-431 (D) and MCF7 (E) whole cell lysates.



HPRT (F-12): sc-373979. Western blot analysis of HPRT expression in A549 (A) and MIA PaCa-2 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.