

CLIM-1/2 (C-4): sc-373985

BACKGROUND

CLIM-1/2 (LIM domain-binding factor-2, LDB1, CLP36, ELFIN, CLP-36, enigma homolog) is a LIM domain-containing protein. LIM domain-containing proteins are capable of binding to a variety of transcription factors. They are likely to function as enhancers, to bring together diverse transcription factors and form higher order activation complexes, or to block formation of such complexes. Human CLIM-1 is a cytoplasmic protein. CLIM-1 northern blot analysis reveals expression of an approximately 2.0-kb transcript most abundant in heart and skeletal muscle. CLIM-1 interacts with the C-terminal EF hand region of α -Actinin-2 through a LIM domain. CLIM-1 localizes to actin stress fibers in nonmuscle cells and associates with α -actinin via its PDZ-domain.

REFERENCES

1. Sugihara, T.M., et al. 1998. Mouse deformed epidermal autoregulatory factor 1 recruits a LIM domain factor, LMO-4, and CLIM coregulators. *Proc. Natl. Acad. Sci. USA* 95: 15418-15423.
2. Becker, T., et al. 2002. Multiple functions of LIM domain-binding CLIM/NLI/Ldb cofactors during zebrafish development. *Mech. Dev.* 117: 75-85.
3. Vallenius, T., et al. 2002. Clik1: a novel kinase targeted to actin stress fibers by the CLP-36 PDZ-LIM protein. *J. Cell Sci.* 115: 2067-2073.
4. Vallenius, T., et al. 2000. CLP-36 PDZ-LIM protein associates with nonmuscle α -Actinin-1 and α -actinin-4. *J. Biol. Chem.* 275: 11100-11105.
5. Bauer, K., et al. 2000. Human CLP36, a PDZ-domain and LIM-domain protein, binds to α -Actinin-1 and associates with actin filaments and stress fibers in activated platelets and endothelial cells. *Blood* 96: 4236-4245.
6. Cassata, G., et al. 2000. The *Caenorhabditis elegans* Ldb/NLI/Clim orthologue Ldb-1 is required for neuronal function. *Dev. Biol.* 226: 45-56.

CHROMOSOMAL LOCATION

Genetic locus: LDB2 (human) mapping to 4p15.32, LDB1 (human) mapping to 10q24.32; Ldb2 (mouse) mapping to 5 B3, Ldb1 (mouse) mapping to 19 C3.

SOURCE

CLIM-1/2 (C-4) is a mouse monoclonal antibody raised against amino acids 1-300 mapping at the N-terminus of CLIM-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-373985 X, 200 μ g/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

CLIM-1/2 (C-4) is recommended for detection of CLIM-1 and CLIM-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CLIM-1/2 (C-4) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of CLIM-1/2: 36/46 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, CCRF-CEM cell lysate: sc-2225 or Jurkat whole cell lysate: sc-2204.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CLIM-1/2 (C-4): sc-373985. Western blot analysis of CLIM-1/2 expression in IMR-32 (A) and Jurkat (B) nuclear extracts.



CLIM-1/2 (C-4): sc-373985. Western blot analysis of CLIM-1/2 expression in CCRF-CEM whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.