



## PA26 siRNA (h): sc-37420

### BACKGROUND

Cell cycle progression is subject to arrest at G1 and G2 checkpoints in response to DNA damage, presumably to allow time for DNA repair prior to entry into S and M phase, respectively. The p53 tumor suppressor is required for one such G1 checkpoint and functions to upregulate expression of GADD 45 and the mitotic inhibitory protein p21. GADD 45 stimulates DNA excision repair *in vitro* and inhibits entry of cells into S phase, and it apparently acts in concert with GADD 153 in inducing growth arrest. A related DNA-damage inducible gene, GADD 34 synergizes with GADD 45 or GADD 153 in suppressing cell growth. PEG-3 (progression elevated gene-3) shares significant homology with GADD 34 and is inducible by DNA damage. An additional GADD related gene, PA26, is a possible target of p53. Three isoforms of PA26 have been identified as PA26-T1, PA26-T2 and PA26-T3.

### REFERENCES

- Hunter, T. et al. 1994. Cyclins and cancer II: cyclin D and CDK inhibitors come of age. *Cell* 79: 573-582.
- Sherr, C.J. 1994. G1 phase progression: cycling on cue. *Cell* 79: 551-555.
- Smith, M.L., et al. 1994. Interaction of the p53-regulated protein Gadd45 with proliferating cell nuclear antigen. *Science* 266: 1376-1380.
- Gujuluva, C.N., et al. 1994. Effect of UV-irradiation on cell cycle, viability and the expression of p53, gadd153 and gadd45 genes in normal and HPV-immortalized human oral keratinocytes. *Oncogene* 9: 1819-1827.
- Selvakumaran, M., et al. 1994. The novel primary response gene MyD118 and the proto-oncogenes myb, myc, and bcl-2 modulate transforming growth factor b1-induced apoptosis of myeloid leukemia cells. *Mol. Cell. Biol.* 14: 2352-2360.
- Zhan, Q., et al. 1994. The gadd and MyD genes define a novel set of mammalian genes encoding acidic proteins that synergistically suppress cell growth. *Mol. Cell. Biol.* 14: 2361-2371.
- Su, Z.-Z., et al. 1997. Subtraction hybridization identifies a transformation progression associated-gene PEG-3 with sequence homology to a growth arrest and DNA damage-inducible gene. *Proc. Natl. Acad. Sci. USA* 94: 9125-9130.
- Velasco-Miguel, S., et al. 1999. PA26, a novel target of the p53 tumor suppressor and member of the GADD family of DNA damage and growth arrest inducible genes. *Oncogene* 18: 127-137.

### CHROMOSOMAL LOCATION

Genetic locus: SESN1 (human) mapping to 6q21; Sesn1 (mouse) mapping to 10 B2.

### PRODUCT

PA26 siRNA (h) is a target-specific 20-25 nt siRNA designed to knock down gene expression. Each vial contains 3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

PA26 siRNA (h) is recommended for the inhibition of PA26 expression in human cells.

PA26 (C-20): sc-9506 is recommended as a control antibody for Western Blotting (starting dilution 1:100, dilution range 1:100-1:1,000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) protein detection using the recommended secondary reagents listed below.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 60  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238. Semi-quantitative RT-PCR may be performed using RT-PCR Primer: PA26 (h)-PR: sc-37420-PR.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.