SANTA CRUZ BIOTECHNOLOGY, INC.

ACOX2 (D-10): sc-374251



BACKGROUND

ACOX2 (acyl-Coenzyme A oxidase 2), also known as BCOX, BRCOX, THCCox or BRCACOX, is a 681 amino acid protein that localizes to the peroxisome and belongs to the acyl-CoA oxidase family. Expressed in heart, kidney, liver, brain, lung, pancreas, placenta and skeletal muscle, ACOX2 functions as a branched-chain acyl-CoA oxidase that is involved in the degradation of bile acid intermediates and long branched fatty acids in peroxisomes. ACOX2 exists as a heterodimer and uses FAD as a cofactor to catalyze oxidation reactions. Defects in the gene encoding ACOX2 may be associated with Zellweger syndrome, an extremely rare congenital disorder that is characterized by the absence of peroxisomes and usually results in death before six months of age.

REFERENCES

- Vanhove, G.F., et al. 1993. The CoA esters of 2-methyl-branched chain fatty acids and of the bile acid intermediates di- and trihydroxycoprostanic acids are oxidized by one single peroxisomal branched chain acyl-CoA oxidase in human liver and kidney. J. Biol. Chem. 268: 10335-10344.
- Baumgart, E., et al. 1996. Mammalian peroxisomal acyl-CoA oxidases. III. Molecular characterization of human branched chain fatty acyl-CoA oxidase. Ann. N.Y. Acad. Sci. 804: 678-679.
- Baumgart, E., et al. 1996. Molecular characterization of the human peroxisomal branched-chain acyl-CoA oxidase: cDNA cloning, chromosomal assignment, tissue distribution, and evidence for the absence of the protein in Zellweger syndrome. Proc. Natl. Acad. Sci. USA 93: 13748-13753.

CHROMOSOMAL LOCATION

Genetic locus: ACOX2 (human) mapping to 3p14.3.

SOURCE

ACOX2 (D-10) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 9-37 near the N-terminus of ACOX2 of human origin.

PRODUCT

Each vial contains 200 $\mu g\, lg G_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ACOX2 (D-10) is available conjugated to agarose (sc-374251 AC), 500 μ g/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-374251 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374251 PE), fluorescein (sc-374251 FITC), Alexa Fluor[®] 488 (sc-374251 AF488), Alexa Fluor[®] 546 (sc-374251 AF546), Alexa Fluor[®] 594 (sc-374251 AF594) or Alexa Fluor[®] 647 (sc-374251 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-374251 AF680) or Alexa Fluor[®] 790 (sc-374251 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-374251 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

ACOX2 (D-10) is recommended for detection of ACOX2 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ACOX2 siRNA (h): sc-78421, ACOX2 shRNA Plasmid (h): sc-78421-SH and ACOX2 shRNA (h) Lentiviral Particles: sc-78421-V.

Molecular Weight of ACOX2: 77 kDa.

Positive Controls: ACOX2 (h): 293T Lysate: sc-116061 or human liver extract: sc-363766.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA





ACOX2 (D-10): sc-374251. Western blot analysis of ACOX2 expression in non-transfected: sc-117752 (**A**) and human ACOX2 transfected: sc-116061 (**B**) 293T whole cell lysates.

ACOX2 (D-10): sc-374251. Western blot analysis of ACOX2 expression in human liver tissue extract.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.