

galectin-3 (H-5): sc-374253

BACKGROUND

Galectins are a family of soluble β -galactoside-binding animal lectins that modulate cell-to-cell adhesion and cell-to-extracellular matrix (ECM) interactions and play a role in tumor progression, pre-mRNA splicing and apoptosis. The galectin-3 protein, also known as Mac-2, hMac-2, GALBP, CBP35 or LGALS3, contains a single carbohydrate binding domain, which binds galactose-containing glycoconjugates. Galectin-3 is expressed in colonic and intestinal epithelium, inflammatory macrophages, papillary and follicular carcinomas, neoplastic astrocytes and some B and T lymphocytes. Upregulated expression of galectin-3 is involved in cancer progression and metastasis. Galectin-3 mediates the endocytosis of β 1 Integrins in a lactose-dependent manner and is associated with thyroid malignancy and Crohn's disease. It may also be used as a marker for diagnosing cases involving Hurthle cell adenomas and carcinomas.

CHROMOSOMAL LOCATION

Genetic locus: LGALS3 (human) mapping to 14q22.3; Lgals3 (mouse) mapping to 14 C1.

SOURCE

galectin-3 (H-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-31 at the N-terminus of galectin-3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374253 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

galectin-3 (H-5) is recommended for detection of galectin-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for galectin-3 siRNA (h): sc-155994, galectin-3 siRNA (m): sc-35443, galectin-3 shRNA Plasmid (h): sc-155994-SH, galectin-3 shRNA Plasmid (m): sc-35443-SH, galectin-3 shRNA (h) Lentiviral Particles: sc-155994-V and galectin-3 shRNA (m) Lentiviral Particles: sc-35443-V.


Molecular Weight of galectin-3: 31 kDa.

Positive Controls: SW480 nuclear extract: sc-2155, A-431 nuclear extract: sc-2122 or MCF7 nuclear extract: sc-2149.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



galectin-3 (H-5): sc-374253. Western blot analysis of galectin-3 expression in T98G (A) and RAW 264.7 (B) whole cell lysates and SW480 (C), A-431 (D), MCF7 (E) and WEHI-231 (F) nuclear extracts.



galectin-3 (H-5): sc-374253. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human colon tissue showing cytoplasmic and nuclear staining of glandular cells and cytoplasmic staining of endothelial cells (B).

SELECT PRODUCT CITATIONS

1. Song, M., et al. 2017. Aluminum trichloride inhibits the rat osteoblasts mineralization *in vitro*. Biol. Trace Elem. Res. 175: 186-193.
2. de Freitas Pedrosa, L., et al. 2020. The acid and neutral fractions of pectins isolated from ripe and overripe papayas differentially affect galectin-3 inhibition and colon cancer cell growth. Int. J. Biol. Macromol. 164: 2681-2690.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **galectin-3 (B2C10): sc-32790** for galectin-3 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.