

# DSCD75 (A-8): sc-374256

## BACKGROUND

Mesenchymal stem cell protein DSCD75, also known as UPF0670 protein C8orf55, is a 208 amino acid secreted protein that belongs to the UPF0670 family. The gene encoding DSCD75 maps to human chromosome 8, which consists of nearly 146 million base pairs, encodes over 800 genes and is associated with a variety of diseases and malignancies. Schizophrenia, bipolar disorder, Trisomy 8, Pfeiffer syndrome, congenital hypothyroidism, Waardenburg syndrome and some leukemias and lymphomas are thought to occur as a result of defects in specific genes that map to chromosome 8.

## REFERENCES

1. Kashino, G., et al. 2001. Preferential expression of an intact WRN gene in Werner syndrome cell lines in which a normal chromosome 8 has been introduced. *Biochem. Biophys. Res. Commun.* 289: 111-115.
2. Selicorni, A., et al. 2002. Cytogenetic mapping of a novel locus for type II Waardenburg syndrome. *Hum. Genet.* 110: 64-67.
3. McQueen, M.B., et al. 2005. Combined analysis from eleven linkage studies of bipolar disorder provides strong evidence of susceptibility loci on chromosomes 6q and 8q. *Am. J. Hum. Genet.* 77: 582-595.
4. Agrelo, R., et al. 2006. Epigenetic inactivation of the premature aging Werner syndrome gene in human cancer. *Proc. Natl. Acad. Sci. USA* 103: 8822-8827.
5. Mossafa, H., et al. 2006. Non-Hodgkin's lymphomas with Burkitt-like cells are associated with c-Myc amplification and poor prognosis. *Leuk. Lymphoma* 47: 1885-1893.
6. Burkard, T.R., et al. 2011. Initial characterization of the human central proteome. *BMC Syst. Biol.* 5: 17.

## CHROMOSOMAL LOCATION

Genetic locus: THEM6 (human) mapping to 8q24.3; Them6 (mouse) mapping to 15 D3.

## SOURCE

DSCD75 (A-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 55-85 within an internal region of DSCD75 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>3</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-374256 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

DSCD75 (A-8) is recommended for detection of DSCD75 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DSCD75 siRNA (h): sc-77625, DSCD75 siRNA (m): sc-143174, DSCD75 shRNA Plasmid (h): sc-77625-SH, DSCD75 shRNA Plasmid (m): sc-143174-SH, DSCD75 shRNA (h) Lentiviral Particles: sc-77625-V and DSCD75 shRNA (m) Lentiviral Particles: sc-143174-V.


Molecular Weight of DSCD75: 24 kDa.

Positive Controls: mouse brain extract: sc-2253, human cerebellum extract: sc-516706 or DSCD75 (h): 293T Lysate: sc-110634.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



DSCD75 (A-8): sc-374256. Western blot analysis of DSCD75 expression in non-transfected: sc-117752 (A) and human DSCD75 transfected: sc-110634 (B) 293T whole cell lysates.



DSCD75 (A-8): sc-374256. Western blot analysis of DSCD75 expression in human cerebellum tissue extract.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.