

CD3- ζ (D-4): sc-374281

BACKGROUND

The T cell antigen receptor (TCR) recognizes foreign antigens and translates such recognition events into intracellular signals that elicit a change in the cell from a dormant to an activated state. Much of this signaling process can be attributed to a multisubunit complex of proteins that associates directly with the TCR. This complex has been designated CD3 (cluster of differentiation 3). It is composed of five invariant polypeptide chains that associate to form three dimers: a heterodimer of γ and ϵ chains ($\gamma\epsilon$), a heterodimer of δ and ϵ chains ($\delta\epsilon$) and a homodimer of two ζ chains ($\zeta\zeta$) or a heterodimer of ζ and η chains ($\zeta\eta$). The ζ and η chains are encoded by the same gene but differ in their carboxyl-terminal ends due to an alternative splicing event. The γ , ϵ and δ chains each contain a single copy of a conserved immunoreceptor tyrosine-based activation motif (ITAM). In contrast, the ζ chain contains three consecutive copies of the same motif. Phosphorylated ITAMs act as docking sites for protein kinases such as ZAP-70 and Syk and are also capable of regulating their kinase activity. The crystal structure of the ZAP-70 SH2 domains bound to the ζ chain ITAMs has been solved.

REFERENCE

1. Exley, M., et al. 1991. Structure, assembly and intracellular transport of the T cell receptor for antigen. *Sem. Immunol.* 3: 283-297.
2. Weiss, A., et al. 1991. Signal transduction by the T cell antigen receptor. *Sem. Immunol.* 3: 313-324.
3. Chan, A.C., et al. 1994. The role of protein tyrosine kinases and protein tyrosine phosphatases in cell antigen receptor signal transduction. *Sem. Immunol.* 12: 555-592.

CHROMOSOMAL LOCATION

Genetic locus: CD247 (human) mapping to 1q24.2; Cd247 (mouse) mapping to 1 H2.3.

SOURCE

CD3- ζ (D-4) is a mouse monoclonal antibody raised against amino acids 1-163 representing full length CD3- ζ of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CD3- ζ (D-4) is available conjugated to agarose (sc-374281 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-374281 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374281 PE), fluorescein (sc-374281 FITC), Alexa Fluor[®] 488 (sc-374281 AF488), Alexa Fluor[®] 546 (sc-374281 AF546), Alexa Fluor[®] 594 (sc-374281 AF594) or Alexa Fluor[®] 647 (sc-374281 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-374281 AF680) or Alexa Fluor[®] 790 (sc-374281 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS


CD3- ζ (D-4) is recommended for detection of CD3- ζ of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CD3- ζ siRNA (h): sc-29245, CD3- ζ/η siRNA (m): sc-42754, CD3- ζ shRNA Plasmid (h): sc-29245-SH, CD3- ζ/η shRNA Plasmid (m): sc-42754-SH, CD3- ζ shRNA (h) Lentiviral Particles: sc-29245-V and CD3- ζ/η shRNA (m) Lentiviral Particles: sc-42754-V.


Molecular Weight of CD3- ζ : 22 kDa.

Positive Controls: BYDP whole cell lysate: sc-364368, MOLT-4 cell lysate: sc-2233 or Jurkat whole cell lysate: sc-2204.

DATA



CD3- ζ (D-4): sc-374281. Western blot analysis of CD3- ζ expression in Jurkat (A), SUP-T1 (B), MOLT-4 (C) and BYDP (D) whole cell lysates.



CD3- ζ (D-4): sc-374281. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lymph node tissue showing germinal centers (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human spleen tissue showing membrane and cytoplasmic staining of cells in white pulp and cells in red pulp (B).

SELECT PRODUCT CITATIONS

1. Esposito, L., et al. 2014. Investigation of soluble and transmembrane CTLA-4 isoforms in serum and microvesicles. *J. Immunol.* 193: 889-900.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.