

Integrin β 1 (A-4): sc-374429

BACKGROUND

Integrins are heterodimers composed of non-covalently associated transmembrane α and β subunits. The 16 α and 8 β subunits heterodimerize to produce more than 20 different receptors. Most integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, collagen and Vitronectin. Certain integrins can also bind to soluble ligands such as Fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis.

CHROMOSOMAL LOCATION

Genetic locus: ITGB1 (human) mapping to 10p11.22; Itgb1 (mouse) mapping to 8 E2.

SOURCE

Integrin β 1 (A-4) is a mouse monoclonal antibody raised against amino acids 375-480 mapping within an extracellular domain of Integrin β 1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Integrin β 1 (A-4) is available conjugated to agarose (sc-374429 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-374429 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374429 PE), fluorescein (sc-374429 FITC), Alexa Fluor[®] 488 (sc-374429 AF488), Alexa Fluor[®] 546 (sc-374429 AF546), Alexa Fluor[®] 594 (sc-374429 AF594) or Alexa Fluor[®] 647 (sc-374429 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-374429 AF680) or Alexa Fluor[®] 790 (sc-374429 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

Integrin β 1 (A-4) is recommended for detection of Integrin β 1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Integrin β 1 siRNA (h): sc-35674, Integrin β 1 siRNA (m): sc-35675, Integrin β 1 siRNA (r): sc-72028, Integrin β 1 shRNA Plasmid (h): sc-35674-SH, Integrin β 1 shRNA Plasmid (m): sc-35675-SH, Integrin β 1 shRNA Plasmid (r): sc-72028-SH, Integrin β 1 shRNA (h) Lentiviral Particles: sc-35674-V, Integrin β 1 shRNA (m) Lentiviral Particles: sc-35675-V and Integrin β 1 shRNA (r) Lentiviral Particles: sc-72028-V.


Molecular Weight of Integrin β 1: 138 kDa.

Positive Controls: SK-N-SH cell lysate: sc-2410 or F9 cell lysate: sc-2245.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Integrin β 1 (A-4): sc-374429. Western blot analysis of Integrin β 1 expression in A549 (A), NIH/3T3 (B), C6 (C) and RT-4 (D) whole cell lysates.



Integrin β 1 (A-4): sc-374429. Western blot analysis of Integrin β 1 expression in U-87 MG (A), SK-N-SH (B) and F9 (C) whole cell lysates.

SELECT PRODUCT CITATIONS

- Shi, Z., et al. 2010. The neuroprotective effect of Batch-2, an aqueous extract from cat's claw (*Uncaria tomentosa*) on 6-OHDA-induced SH-SY5Y cell damage. *Prog. Biochem. Biophys.* 37: 769-778.
- Bartolomé, R.A., et al. 2014. Cadherin-17 interacts with α 2 β 1 Integrin to regulate cell proliferation and adhesion in colorectal cancer cells causing liver metastasis. *Oncogene* 33: 1658-1669.
- Shen, Y., et al. 2015. Effect of surface chemistry on the integrin induced pathway in regulating vascular endothelial cells migration. *Colloids Surf. B, Biointerfaces* 126: 188-197.
- Huang, L.H., et al. 2016. Myeloid acyl-CoA:cholesterol acyltransferase 1 deficiency reduces lesion macrophage content and suppresses atherosclerosis progression. *J. Biol. Chem.* 291: 6232-6244.
- Morelle, W., et al. 2017. Galactose supplementation in patients with TMEM165-CDG rescues the glycosylation defects. *J. Clin. Endocrinol. Metab.* 102: 1375-1386.
- Kim, Y.R., et al. 2018. Integrin α 6 as an invasiveness marker for hepatitis B viral X-driven hepatocellular carcinoma. *Cancer Biomark.* 23: 135-144.
- Husari, A., et al. 2019. On the relationship of YAP and FAK in hMSCs and osteosarcoma cells: discrimination of FAK modulation by nuclear YAP depletion or YAP silencing. *Cell. Signal.* 63: 109382.
- Li, X., et al. 2020. Therapeutic ultrasound combined with microbubbles improves atherosclerotic plaque stability by selectively destroying the intraplaque neovasculature. *Theranostics* 10: 2522-2537.
- Iwai, M., et al. 2021. Cancer-associated fibroblast migration in non-small cell lung cancers is modulated by increased Integrin α 11 expression. *Mol. Oncol.* 15: 1507-1527.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA