

Cdx4 (H-4): sc-374471



The Power to Question

BACKGROUND

Cdx1, Cdx2 and Cdx4 are members of the caudal-type homeobox family of genes, which are homologs of the *Drosophila* "caudal" gene required for anterior-posterior regional identity. The proteins encoded by these genes are transcription factors which play an important role in development by regulating the expression of Hox genes. Hox genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. Cdx4 is a major positive regulator of the expression of all Hox family members. Due to its critical role as a regulator, Cdx4 is a direct target of the canonical Wnt pathway. The loss of Cdx4 can result in the development of an expanded hindbrain, while the overexpression of Cdx4 may cause the hindbrain to lose its distinct segmental features and resemble the spinal cord.

REFERENCES

1. Gamer, L.W., et al. 1993. Murine Cdx-4 bears striking similarities to the *Drosophila* caudal gene in its homeodomain sequence and early expression pattern. *Mech. Dev.* 43: 71-81.
2. Horn, J.M., et al. 1995. A member of the caudal family of homeobox genes maps to the X-inactivation centre region of the mouse and human X chromosomes. *Hum. Mol. Genet.* 4: 1041-1047.
3. Bonner, C.A., et al. 1995. Isolation, characterization, and precise physical localization of human CDX1, a caudal-type homeobox gene. *Genomics* 28: 206-211.
4. Gaunt, S.J., et al. 2005. Cdx4/lacZ and Cdx2/lacZ protein gradients formed by decay during gastrulation in the mouse. *Int. J. Dev. Biol.* 49: 901-908.
5. Pilon, N., et al. 2006. Cdx4 is a direct target of the canonical Wnt pathway. *Dev. Biol.* 289: 55-63.

CHROMOSOMAL LOCATION

Genetic locus: CDX4 (human) mapping to Xq13.2; Cdx4 (mouse) mapping to X D.

SOURCE

Cdx4 (H-4) is a mouse monoclonal antibody raised against amino acids 1-168 mapping at the N-terminus of Cdx4 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-374471 X, 200 µg/0.1 ml.

Cdx4 (H-4) is available conjugated to agarose (sc-374471 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-374471 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-374471 PE), fluorescein (sc-374471 FITC), Alexa Fluor® 488 (sc-374471 AF488), Alexa Fluor® 546 (sc-374471 AF546), Alexa Fluor® 594 (sc-374471 AF594) or Alexa Fluor® 647 (sc-374471 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-374471 AF680) or Alexa Fluor® 790 (sc-374471 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

Cdx4 (H-4) is recommended for detection of Cdx4 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Cdx4 siRNA (h): sc-72316, Cdx4 siRNA (m): sc-72317, Cdx4 shRNA Plasmid (h): sc-72316-SH, Cdx4 shRNA Plasmid (m): sc-72317-SH, Cdx4 shRNA (h) Lentiviral Particles: sc-72316-V and Cdx4 shRNA (m) Lentiviral Particles: sc-72317-V.

Cdx4 (H-4) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Cdx4: 30 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Hep G2 cell lysate: sc-2227 or SK-N-MC nuclear extract: sc-2154.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Cdx4 (H-4): sc-374471. Western blot analysis of Cdx4 expression in SK-N-MC nuclear extract (A) and SK-N-SH (B), SH-SY5Y (C), IMR-32 (D) and Neuro-2A (E) whole cell lysates.



Cdx4 (H-4): sc-374471. Fluorescent western blot analysis of Cdx4 expression in HeLa (A), c4 (B), Hep G2 (C) and SK-N-MC (D) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgGκ BP-CFL 647: sc-516179.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA