



CENP-H siRNA (m): sc-37566

BACKGROUND

A replicated chromosome includes two kinetochores that control chromosome segregation during mitosis. Both centromere proteins CENP-B and CENP-H are contained in the centromeric heterochromatin between kinetochores, and are involved in maintaining sister chromatid cohesion. The highly dispersed CENP-B promotes and maintains the joining of DNA satellites in the centromere. CENP-B targets centromeric α -DNA and protects it from digestion by nucleases as well as preventing DNase or restriction enzyme digestion from affecting the morphology of centromeres. CENP-H contains a coiled-coil structure and a nuclear localization signal. CENP-H is specifically and constitutively localized to kinetochores and plays a role in the organization and function of kinetochores throughout the cell cycle.

REFERENCES

1. Cooke, C.A., et al. 1990. CENP-B: a major human centromere protein located beneath the kinetochore. *J. Cell Biol.* 110: 1475-1488.
2. Rieder, C.L., et al. 1998. The vertebrate cell kinetochore and its roles during mitosis. *Trends Cell Biol.* 8: 310-318.
3. Barbosa-Cisneros, O., et al. 1998. Localization of the centromere protein CENP-B using scleroderma sera and evidence for a role in centromere survival. *Rev. Rhum. Engl. Ed* 65: 15-20.
4. Sugata, N., et al. 1999. Characterization of a novel kinetochore protein, CENP-H. *J. Biol. Chem.* 274: 27343-27346.
5. Choo, K.H. 2000. Centromerization. *Trends Cell Biol.* 10: 182-188.
6. Izuta, H., et al. 2006. Comprehensive analysis of the ICEN (interphase centromere complex) components enriched in the CENP-A chromatin of human cells. *Genes Cells* 11: 673-684.
7. Orthaus, S., et al. 2006. RNAi knockdown of human kinetochore protein CENP-H. *Biochem. Biophys. Res. Commun.* 348: 36-46.
8. Shigeishi, H., et al. 2006. Increased expression of CENP-H gene in human oral squamous cell carcinomas harboring high-proliferative activity. *Oncol. Rep.* 16: 1071-1075.

CHROMOSOMAL LOCATION

Genetic locus: Cenph (mouse) mapping to 13 D1.

PRODUCT

CENP-H siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CENP-H shRNA Plasmid (m): sc-37566-SH and CENP-H shRNA (m) Lentiviral Particles: sc-37566-V as alternate gene silencing products.

For independent verification of CENP-H (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-37566A, sc-37566B and sc-37566C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CENP-H siRNA (m) is recommended for the inhibition of CENP-H expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CENP-H gene expression knockdown using RT-PCR Primer: CENP-H (m)-PR: sc-37566-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.