



cyclin T2a/b siRNA (m): sc-37602

BACKGROUND

Cyclin T1 was identified as a partner for Cdk9, an RNA polymerase II (RNAPII) transcription elongation factor. cyclin T1 interacts with the transactivation domain of the HIV-1 Tat protein. The interaction of Tat with cyclin T1 enhances the affinity of Tat for the viral TAR RNA stem-loop structure, suggesting that Tat can recruit cyclin T1/Cdk9 to RNAPII through cooperative binding to TAR. The human positive transcription elongation factor b (P-TEFb) consists of a cyclin dependent kinase, Cdk9, paired with a cyclin T. Cdk9 may be paired with either cyclin T1 or cyclin T2, in a mutually exclusive manner. Two forms of cyclin T2, T2a and T2b, are due to alternative splicing. The binding of Tat to TAR was shown to be facilitated by human cyclin T1, but not by cyclins T2a or T2b. cyclin T2 binds to Cdk9 but not to Tat, and cyclin T2 can inhibit cyclin T1-mediated Tat activity.

REFERENCES

1. Herrmann, C.H., et al. 1995. Lentivirus Tat proteins specifically associate with a cellular protein kinase, TAK, that hyperphosphorylates the carboxyl-terminal domain of the large subunit of RNA polymerase II: candidate for a Tat cofactor. *J. Virol.* 69: 1612-1620.
2. Yang, X., et al. 1997. TAK, an HIV Tat-associated kinase, is a member of the cyclin-dependent family of protein kinases and is induced by activation of peripheral blood lymphocytes and differentiation of promonocytic cell lines. *Proc. Natl. Acad. Sci. USA* 94: 12331-12336.
3. Wei, P., et al. 1998. A novel Cdk9-associated C-type cyclin interacts directly with HIV-1 Tat and mediates its high-affinity, loop-specific binding to TAR RNA. *Cell* 92: 451-462.
4. Peng, J., et al. 1998. Identification of multiple cyclin subunits of human P-TEFb. *Genes Dev.* 12: 755-762.
5. Wimmer, J., et al. 1999. Interactions between Tat and TAR and human immunodeficiency virus replication are facilitated by human cyclin T1 but not cyclins T2a or T2b. *Virology* 255: 182-189.
6. Napolitano, G., et al. 1999. The Cdk9-associated cyclins T1 and T2 exert opposite effects on HIV-1 Tat activity. *AIDS* 13: 1453-1459.
7. Cottone, G. et al. 2006. Pkn is a novel partner of cyclin T2a in muscle differentiation. *J. Cell. Physiol.* 207: 232-237.
8. Liou, L.Y., et al. 2006. Cyclin T1 but not cyclin T2a is induced by a post-transcriptional mechanism in PAMP-activated monocyte-derived macrophages. *J. Leukoc. Biol.* 79: 388-396.

CHROMOSOMAL LOCATION

Genetic locus: Ccnt2 (mouse) mapping to 1 E3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

cyclin T2a/b siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see cyclin T2a/b shRNA Plasmid (m): sc-37602-SH and cyclin T2a/b shRNA (m) Lentiviral Particles: sc-37602-V as alternate gene silencing products.

For independent verification of cyclin T2a/b (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-37602A, sc-37602B and sc-37602C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

cyclin T2a/b siRNA (m) is recommended for the inhibition of cyclin T2a/b expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor cyclin T2a/b gene expression knockdown using RT-PCR Primer: cyclin T2a/b (m)-PR: sc-37602-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.