

# Rab 4/14 (F-10): sc-376243

## BACKGROUND

The Ras-related superfamily of guanine nucleotide binding proteins, which includes the R-Ras, Rap, Ral/Rec and Rho/Rab subfamilies exhibit 30-60% homology with Ras p21. Accumulating data suggests an important role for Rab proteins, either in endocytosis or in biosynthetic protein transport. The transport of newly synthesized proteins from the endoplasmic reticulum to various stacks of the Golgi complex and to secretory vesicles involves at each stage the movement of carrier vesicles, a process that appears to involve Rab protein function. The possibility that Rab proteins might also direct the exocytosis from secretory vesicles to the plasma membrane is supported by the observation that in yeast, the SEC4 protein, which is 40% homologous to Rab proteins, is associated with secretory vesicles. At least eight members of the Rab subfamily have been identified, each of which is found at a particular stage of a membrane transport pathway.

## SOURCE

Rab 4/14 (F-10) is a mouse monoclonal antibody raised against amino acids 1-213 representing full length Rab 4A of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Rab 4/14 (F-10) is available conjugated to agarose (sc-376243 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376243 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376243 PE), fluorescein (sc-376243 FITC), Alexa Fluor® 488 (sc-376243 AF488), Alexa Fluor® 546 (sc-376243 AF546), Alexa Fluor® 594 (sc-376243 AF594) or Alexa Fluor® 647 (sc-376243 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-376243 AF680) or Alexa Fluor® 790 (sc-376243 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Rab 4/14 (F-10) is recommended for detection of Rab 4A, Rab 4B and Rab 14 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); partially cross reactive with other Rab family members.


Rab 4/14 (F-10) is also recommended for detection of Rab 4A, Rab 4B and Rab 14 in additional species, including equine.

Positive Controls: Rab 4A (h3): 293T Lysate: sc-173396, MCF7 whole cell lysate: sc-2206 or Hep G2 cell lysate: sc-2227.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



Rab 4/14 (F-10): sc-376243. Western blot analysis of Rab 4/14 expression in non-transfected 293T: sc-117752 (A), human Rab 4A transfected 293T: sc-173396 (B), HeLa (C), MCF7 (D), SH-SY5Y (E) and Hep G2 (F) whole cell lysates.



Rab 4/14 (F-10): sc-376243. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic and membrane localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human appendix tissue showing cytoplasmic staining of glandular and lymphoid cells (B).

## SELECT PRODUCT CITATIONS

- Contreras-García, I.J., et al. 2018. Differential expression of synaptic vesicle protein 2A after status epilepticus and during epilepsy in a lithium-pilocarpine model. *Epilepsy Behav.* 88: 283-294.
- Bian, W., et al. 2023. A spatially defined human Notch receptor interaction network reveals Notch intracellular storage and Ataxin-2-mediated fast recycling. *Cell Rep.* 42: 112819.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.