

PMF-1 (G-9): sc-376269

BACKGROUND

PMF-1 (polyamine-modulated factor 1) is a 205 amino acid protein involved in kinetochore formation. Localized to the nucleus, PMF-1 contains a coiled-coil domain which interacts with the leucine-zipper domain of Nrf2. This interaction regulates the transcription of SSAT, a regulatory enzyme for polyamine catabolism. PMF-1 is also a component of the MIS12 complex, which is required for kinetochore formation and chromosomal alignment and segregation. PMF-1 is expressed at highest levels in skeletal muscle and heart, with moderate expression in liver and kidney. PMF-1 exists as five isoforms produced by alternative splicing.

REFERENCES

- Wang, Y., et al. 1999. Cloning and characterization of human polyamine-modulated factor-1, a transcriptional cofactor that regulates the transcription of the spermidine/spermine N¹-acetyltransferase gene. *J. Biol. Chem.* 274: 22095-22101.
- Wang, Y., et al. 2001. Characterization of the interaction between the transcription factors human polyamine modulated factor (PMF-1) and NF-E2-related factor 2 (Nrf-2) in the transcriptional regulation of the spermidine/spermine N¹-acetyltransferase (SSAT) gene. *Biochem. J.* 355: 45-49.
- Wang, Y., et al. 2002. Polyamine-modulated factor 1 binds to the human homologue of the 7a subunit of the *Arabidopsis* COP9 signalosome: implications in gene expression. *Biochem. J.* 366: 79-86.
- Cheeseman, I.M., et al. 2004. A conserved protein network controls assembly of the outer kinetochore and its ability to sustain tension. *Genes Dev.* 18: 2255-2268.
- Obuse, C., et al. 2004. A conserved Mis12 centromere complex is linked to heterochromatic HP1 and outer kinetochore protein Zwint-1. *Nat. Cell Biol.* 6: 1135-1141.
- Kline, S.L., et al. 2006. The human Mis12 complex is required for kinetochore assembly and proper chromosome segregation. *J. Cell Biol.* 173: 9-17.
- Hyvönen, M.T., et al. 2006. Polyamine-regulated unproductive splicing and translation of spermidine/spermine N¹-acetyltransferase. *RNA* 12: 1569-1582.
- Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 609176. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Aleman, A., et al. 2008. Identification of PMF1 methylation in association with bladder cancer progression. *Clin. Cancer Res.* 14: 8236-8243.

CHROMOSOMAL LOCATION

Genetic locus: PMF1 (human) mapping to 1q22.

SOURCE

PMF-1 (G-9) is a mouse monoclonal antibody raised against amino acids 76-180 mapping within an internal region of PMF-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PMF-1 (G-9) is recommended for detection of PMF-1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PMF-1 siRNA (h): sc-78780, PMF-1 shRNA Plasmid (h): sc-78780-SH and PMF-1 shRNA (h) Lentiviral Particles: sc-78780-V.


Molecular Weight of PMF-1 isoforms: 19/23/24 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, Jurkat nuclear extract: sc-2132 or MCF7 whole cell lysate: sc-2206.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



PMF-1 (G-9): sc-376269. Western blot analysis of PMF-1 expression in Jurkat nuclear extract.



PMF-1 (G-9): sc-376269. Western blot analysis of PMF-1 expression in MCF7 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.