

CENP-B (C-10): sc-376392

BACKGROUND

A replicated chromosome includes two kinetochores that control chromosome segregation during mitosis. Both centromere proteins CENP-B and CENP-H are contained in the centromeric heterochromatin between kinetochores, and are involved in maintaining sister chromatid cohesion. The highly dispersed CENP-B promotes and maintains the joining of DNA satellites in the centromere. CENP-B targets centromeric α -DNA and protects it from digestion by nucleases as well as preventing DNase or restriction enzyme digestion from affecting the morphology of centromeres. CENP-H contains a coiled-coil structure and a nuclear localization signal. CENP-H is specifically and constitutively localized to kinetochores and plays a role in the organization and function of kinetochores throughout the cell cycle.

CHROMOSOMAL LOCATION

Genetic locus: CENPB (human) mapping to 20p13; Cenpb (mouse) mapping to 2 F1.

SOURCE

CENP-B (C-10) is a mouse monoclonal antibody raised against amino acids 535-599 mapping at the C-terminus of CENP-B of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CENP-B (C-10) is available conjugated to agarose (sc-376392 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376392 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376392 PE), fluorescein (sc-376392 FITC), Alexa Fluor® 488 (sc-376392 AF488), Alexa Fluor® 546 (sc-376392 AF546), Alexa Fluor® 594 (sc-376392 AF594) or Alexa Fluor® 647 (sc-376392 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-376392 AF680) or Alexa Fluor® 790 (sc-376392 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CENP-B (C-10) is recommended for detection of CENP-B of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CENP-B siRNA (h): sc-37557, CENP-B siRNA (m): sc-37558, CENP-B shRNA Plasmid (h): sc-37557-SH, CENP-B shRNA Plasmid (m): sc-37558-SH, CENP-B shRNA (h) Lentiviral Particles: sc-37557-V and CENP-B shRNA (m) Lentiviral Particles: sc-37558-V.


Molecular Weight of CENP-B: 80 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, A-431 whole cell lysate: sc-2201 or CENP-B (h): 293T Lysate: sc-116535.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CENP-B (C-10): sc-376392. Western blot analysis of CENP-B expression in non-transfected: sc-117752 (A) and human CENP-B transfected: sc-116535 (B) 293T whole cell lysates.



CENP-B (C-10): sc-376392. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization (A). CENP-B (C-10) Alexa Fluor® 488; sc-376392 AF488. Direct immunofluorescence staining of formalin-fixed SW480 cells showing nuclear bodies localization. Blocked with UltraCruz® Blocking Reagent: sc-516214 (B).

SELECT PRODUCT CITATIONS

- Gentili, M., et al. 2019. The N-terminal domain of cGAS determines preferential association with centromeric DNA and innate immune activation in the nucleus. *Cell Rep.* 26: 2377-2393.e13.
- Traynor, S., et al. 2019. Remodeling and destabilization of chromosome 1 pericentromeric heterochromatin by SSX proteins. *Nucleic Acids Res.* 47: 6668-6684.
- Chunduri, N.K., et al. 2021. Systems approaches identify the consequences of monosomy in somatic human cells. *Nat. Commun.* 12: 5576.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA