

STK33 (G-11): sc-376498

BACKGROUND

The phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including cell division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/threonine (Ser/Thr) protein kinases. STK33 (serine/threonine kinase 33) is a 514 amino acid protein that belongs to the CaMK (calcium/calmodulin dependent kinase) subfamily of structurally related serine/threonine kinases. Widely expressed at low levels with predominant expression in testis, lung, retina and fetal organs such as brain, heart and spinal cord, STK33 contains one protein kinase domain and functions as a Ser/Thr protein kinase with a possible role in spermatogenesis. The gene encoding STK33 lies within a region on chromosome 11 that has been associated with a variety of defects, including Long QT syndrome, T-cell leukemia, Beckwith-Wiedemann syndrome, Usher syndrome 1C and various other malignancies.

REFERENCES

- Amid, C., et al. 2001. Comparative genomic sequencing reveals a strikingly similar architecture of a conserved syntenic region on human chromosome 11p15.3 (including gene ST5) and mouse chromosome 7. *Cytogenet. Cell Genet.* 93: 284-290.
- Mujica, A.O., et al. 2001. A novel serine/threonine kinase gene, STK33, on human chromosome 11p15.3. *Gene* 280: 175-181.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607670. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Guo, L., et al. 2003. Molecular cloning and characterization of a novel human kinase gene, PDIK1L. *J. Genet.* 82: 27-32.
- Mujica, A.O., et al. 2005. Differential expression pattern of the novel serine/threonine kinase, STK33, in mice and men. *FEBS J.* 272: 4884-4898.

CHROMOSOMAL LOCATION

Genetic locus: STK33 (human) mapping to 11p15.4.

SOURCE

STK33 (G-11) is a mouse monoclonal antibody raised against amino acids 161-236 mapping within an internal region of STK33 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STK33 (G-11) is available conjugated to agarose (sc-376498 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376498 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376498 PE), fluorescein (sc-376498 FITC), Alexa Fluor® 488 (sc-376498 AF488), Alexa Fluor® 546 (sc-376498 AF546), Alexa Fluor® 594 (sc-376498 AF594) or Alexa Fluor® 647 (sc-376498 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-376498 AF680) or Alexa Fluor® 790 (sc-376498 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

STK33 (G-11) is recommended for detection of STK33 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for STK33 siRNA (h): sc-96894, STK33 shRNA Plasmid (h): sc-96894-SH and STK33 shRNA (h) Lentiviral Particles: sc-96894-V.


Molecular Weight of STK33: 53 kDa.

Positive Controls: U-251-MG whole cell lysate: sc-364176, HEK293 whole cell lysate: sc-45136 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



STK33 (G-11) Alexa Fluor® 647: sc-376498 AF647. Direct fluorescent western blot analysis of STK33 expression in U-251-MG (A), HEK293 (B), Daudi (C) and U266 (D) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214.



STK33 (G-11): sc-376498. Western blot analysis of STK33 expression in HeLa whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA