

Lipin-1 (B-12): sc-376874

BACKGROUND

The Lipin family of nuclear proteins contains three members: Lipin-1, Lipin-2 and Lipin-3, all of which contain a nuclear signal sequence, a highly conserved amino-terminal (NLIP) domain, and a carboxy-terminal (CLIP) domain. Lipin-1 is crucial for normal adipose tissue development and metabolism. Lipin-1 selectively activates a subset of PGC-1 α target pathways, including fatty acid oxidation and mitochondrial oxidative phosphorylation by inducing expression of the nuclear receptor PPAR α . Lipin-1 also inactivates the lipogenic program and suppresses circulating lipid levels. An abundance of Lipin-1 promotes fat accumulation and Insulin sensitivity, whereas a deficiency in Lipin-1 may deter normal adipose tissue development, resulting in Insulin resistance and lipodystrophy, a heterogeneous group of disorders characterized by loss of body fat, fatty liver, hypertriglyceridemia and Insulin resistance.

CHROMOSOMAL LOCATION

Genetic locus: LPIN1 (human) mapping to 2p25.1; Lpin1 (mouse) mapping to 12 A1.1.

SOURCE

Lipin-1 (B-12) is a mouse monoclonal antibody raised against amino acids 261-380 mapping within an internal region of Lipin-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Lipin-1 (B-12) is available conjugated to agarose (sc-376874 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376874 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376874 PE), fluorescein (sc-376874 FITC), Alexa Fluor[®] 488 (sc-376874 AF488), Alexa Fluor[®] 546 (sc-376874 AF546), Alexa Fluor[®] 594 (sc-376874 AF594) or Alexa Fluor[®] 647 (sc-376874 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-376874 AF680) or Alexa Fluor[®] 790 (sc-376874 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Lipin-1 (B-12) is recommended for detection of Lipin-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Lipin-1 siRNA (h): sc-60940, Lipin-1 siRNA (m): sc-60941, Lipin-1 shRNA Plasmid (h): sc-60940-SH, Lipin-1 shRNA Plasmid (m): sc-60941-SH, Lipin-1 shRNA (h) Lentiviral Particles: sc-60940-V and Lipin-1 shRNA (m) Lentiviral Particles: sc-60941-V.


Molecular Weight of Lipin-1: 102 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204 or AN3 CA cell lysate: sc-24662.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



Lipin-1 (B-12): sc-376874. Western blot analysis of Lipin-1 expression in Jurkat whole cell lysate.



Lipin-1 (B-12): sc-376874. Western blot analysis of Lipin-1 expression in AN3 CA whole cell lysate.

SELECT PRODUCT CITATIONS

- Shimizu, K., et al. 2017. The SCF β -TRCP E3 ubiquitin ligase complex targets Lipin-1 for ubiquitination and degradation to promote hepatic lipogenesis. *Sci. Signal.* 10: eaah4117.
- Mingorance, L., et al. 2018. Host phosphatidic acid phosphatase Lipin-1 is rate limiting for functional hepatitis C virus replicase complex formation. *PLoS Pathog.* 14: e1007284.
- Li, T.Y., et al. 2018. Tip60-mediated lipin 1 acetylation and ER translocation determine triacylglycerol synthesis rate. *Nat. Commun.* 9: 1916.
- Castro, V., et al. 2019. Differential roles of Lipin-1 and Lipin-2 in the Hepatitis C virus replication cycle. *Cells* 8: 1456.
- Rashid, T., et al. 2019. Lipin-1 deficiency causes sarcoplasmic reticulum stress and chaperone-responsive myopathy. *EMBO J.* 38: e99576.

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.