SANTA CRUZ BIOTECHNOLOGY, INC.

SIRP-α (C-7): sc-376884



BACKGROUND

SIRPs (signal-regulatory proteins) are a family of transmembrane glycoproteins that were identified by their association with the Src homology 2 domaincontaining protein-tyrosine phosphatase SHP-2 in response to Insulin. The SIRP family negatively regulates the PI 3-kinase pathway, which may diminish EGFR-mediated motility and survival phenotypes that contribute to transformation of certain cell types. SIRP- α 1 is a transmembrane protein which contains an extracellular portion with three immunoglobulin-like structures and a cytoplasmic region with four potential tyrosine phosphorylation sites. SIRP- α (also known as SIRP- α 1, SIRP- α 2 or SIRP- α 3) is a substrate for activated receptor tyrosine kinases. In its tyrosine phosphorylated form, SIRP- α binds to SH-PTP2 through SH2 interactions and acts as an SH-PTP2 substrate. SIRP- α has been shown to have negative regulatory effects on cellular responses induced by growth factors, oncogenes and Insulin. SIRP-B1 shares extensive sequence homology with SIRP- α in its extracellular portion but lacks the cytoplasmic portion. SIRP-y, originally designated SIRP-B2 (SIRP-B2, CD172y) has unique characteristics from both the α and β versions. SIRP- γ is expressed on the majority of T cells and a proportion of B cells. CD47 associates with SIRP-y, and this interaction signals unidirectionally only.

CHROMOSOMAL LOCATION

Genetic locus: SIRPA (human) mapping to 20p13; Sirpa (mouse) mapping to 2 F1.

SOURCE

SIRP- α (C-7) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 475-503 at the C-terminus of SIRP- α of human origin.

PRODUCT

Each vial contains 200 μg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

SIRP- α (C-7) is available conjugated to agarose (sc-376884 AC), 500 µg/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-376884 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376884 PE), fluorescein (sc-376884 FITC), Alexa Fluor[®] 488 (sc-376884 AF488), Alexa Fluor[®] 546 (sc-376884 AF546), Alexa Fluor[®] 594 (sc-376884 AF594) or Alexa Fluor[®] 647 (sc-376884 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-376884 AF680) or Alexa Fluor[®] 790 (sc-376884 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-376884 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

SIRP- α (C-7) is recommended for detection of SIRP- α of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SIRP- α siRNA (h): sc-44106, SIRP- α siRNA (m): sc-36493, SIRP- α siRNA (r): sc-270499, SIRP- α shRNA Plasmid (h): sc-44106-SH, SIRP- α shRNA Plasmid (m): sc-36493-SH, SIRP- α shRNA Plasmid (r): sc-270499-SH, SIRP- α shRNA (h) Lentiviral Particles: sc-44106-V, SIRP- α shRNA (m) Lentiviral Particles: sc-36493-V and SIRP- α shRNA (r) Lentiviral Particles: sc-270499-V.

Molecular Weight of unglycosylated SIRP-a: 65 kDa.

Molecular Weight of glycosylated SIRP- α : 100-150 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, THP-1 cell lysate: sc-2238 or SIRP- α (h): 293T Lysate: sc-159295.

DATA





SIRP- α (C-7): sc-376884. Western blot analysis of SIRP- α expression in non-transfected: sc-117752 (**A**) and human SIRP- α transfected: sc-159295 (**B**) 293T whole cell lysates.

SIRP- α (C-7): sc-376884. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cerebellum tissue showing cytoplasmic staining of Purkinje cells.

SELECT PRODUCT CITATIONS

- Deshpande, R.P., et al. 2017. SIRP-α protein downregulates in human astrocytoma: presumptive involvement of Hsa-miR-520d-5p and Hsa-miR-520d-3p. Mol. Neurobiol. 54: 8162-8169.
- Ghimire, K., et al. 2019. Deficiency in SIRP-α cytoplasmic recruitment confers protection from acute kidney injury. FASEB J. 33: 11528-11540.
- Mun, S., et al. 2022. Transcriptome profile of membrane and extracellular matrix components in ligament-fibroblastic progenitors and cementoblasts differentiated from human periodontal ligament cells. Genes 13: 659.
- 4. Lu, J., et al. 2023. Reprogramming of TAMs via the STAT3/CD47-SIRP α axis promotes acquired resistance to EGFR-TKIs in lung cancer. Cancer Lett. 564: 216205.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA