

# RKIP (H-10): sc-376925

## BACKGROUND

Raf kinase inhibitor protein (RKIP) is a cytosolic protein that was initially characterized as a phosphatidylethanolamine-binding protein (PBP) expressed in brain tissue and secreted from testis fluid. In addition, RKIP was identified by yeast two-hybrid screening of human T cell libraries directed at identifying proteins that associate with the BXB kinase domain of the serine/threonine kinase, Raf-1. Subsequent *in vitro* and *in vivo* studies indicate that RKIP binds to both the active and inactive forms of Raf-1 and thereby regulates the signaling cascade of the MAP kinase pathway. The specific association of RKIP with kinase-active Raf-1 competitively inhibits the binding and activation of the Raf-1 substrate MEK. RKIP, in turn, affects downstream MAP kinase signaling by decreasing the activation of MEK effector proteins, including ERK1 and ERK2, and the subsequent induction of AP-1 mediated transcription.

## REFERENCES

1. Perry, A.C., et al. 1994. Sequence analysis of a mammalian phospholipid-binding protein from testis and epididymis and its distribution between spermatozoa and extracellular secretions. *Biochem. J.* 301: 235-242.
2. Minden, A., et al. 1994. Differential activation of ERK and JNK mitogen-activated protein kinases by Raf-1 and MEKK. *Science* 266: 1719-1723.
3. Tohdoh, N., et al. 1995. Sequence homology of rat and human HCNP precursor proteins, bovine phosphatidylethanol-amine-binding protein and rat 23 kDa protein associated with the opioid-binding protein. *Brain Res. Mol. Brain Res.* 30: 381-384.

## CHROMOSOMAL LOCATION

Genetic locus: PEBP1 (human) mapping to 12q24.23, XAB2 (human) mapping to 19p13.2; Pebp1 (mouse) mapping to 5 F, Xab2 (mouse) mapping to 8 A1.1.

## SOURCE

RKIP (H-10) is a mouse monoclonal antibody raised against amino acids 1-187 representing full length RKIP of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RKIP (H-10) is available conjugated to agarose (sc-376925 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-376925 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-376925 PE), fluorescein (sc-376925 FITC), Alexa Fluor<sup>®</sup> 488 (sc-376925 AF488), Alexa Fluor<sup>®</sup> 546 (sc-376925 AF546), Alexa Fluor<sup>®</sup> 594 (sc-376925 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-376925 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-376925 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-376925 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

RKIP (H-10) is recommended for detection of RKIP and processed active peptide HCNP of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of RKIP: 23 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, MCF7 whole cell lysate: sc-2206 or c4 whole cell lysate: sc-364186.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



RKIP (H-10): sc-376925. Western blot analysis of RKIP expression in Jurkat (A), MCF7 (B), c4 (C), BC<sub>3</sub>H1 (D), C6 (E) and H19-7/IGF-IR (F) whole cell lysates.



RKIP (H-10): sc-376925. Immunofluorescence staining of methanol-fixed NIH/3T3 cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic and membrane staining of myocytes (B).

## SELECT PRODUCT CITATIONS

1. Ahn, S.Y., et al. 2017. Anti-helminthic niclosamide inhibits Ras-driven oncogenic transformation via activation of GSK-3. *Oncotarget* 8: 31856-31863.
2. Qi, Z.H., et al. 2018. RIPK4/PEBP1 axis promotes pancreatic cancer cell migration and invasion by activating RAF1/MEK/ERK signaling. *Int. J. Oncol.* 52: 1105-1116.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.