

CCDC33 (F-3): sc-377074

BACKGROUND

The coiled-coil domain is a structural motif found in proteins that are involved in a diverse array of biological functions such as the regulation of gene expression, cell division, membrane fusion and drug extrusion and delivery. CCDC33 (coiled-coil domain containing 33), also known as CT61 (cancer/testis antigen 61), is a 958 amino acid protein found primarily in male germ cells. Existing as at least four alternatively spliced isoforms, CCDC33 is implicated in spermatogenesis and contains one C2 domain. CCDC33 is encoded by a gene located on human chromosome 15, which houses over 700 genes and comprises nearly 3% of the human genome. Angelman syndrome, Prader-Willi syndrome, Tay-Sachs disease and Marfan syndrome are all associated with defects in chromosome 15-localized genes.

REFERENCES

- Hurowitz, G.I., et al. 1993. Neuropsychiatric aspects of adult-onset Tay-Sachs disease: two case reports with several new findings. *J. Neuropsychiatry Clin. Neurosci.* 5: 30-36.
- Woolfson, D.N. 2005. The design of coiled-coil structures and assemblies. *Adv. Protein Chem.* 70: 79-112.
- Midla, G.S. 2008. Diagnosis and management of patients with Marfan syndrome. *JAAPA* 21: 21-25.
- Kaczmarek, K., et al. 2009. Ccdc33, a predominantly testis-expressed gene, encodes a putative peroxisomal protein. *Cytogenet. Genome Res.* 126: 243-252.
- Dan, B. 2009. Angelman syndrome: current understanding and research prospects. *Epilepsia* 50: 2331-2339.
- Ferrer-Bolufer, I., et al. 2009. Tyrosinemia type 1 and Angelman syndrome due to paternal uniparental isodisomy 15. *J. Inher. Metab. Dis.* 32: S349-S353.
- Moutevelis, E. and Woolfson, D.N. 2009. A periodic table of coiled-coil protein structures. *J. Mol. Biol.* 385: 726-732.
- Wawrzik, M., et al. 2010. The C15orf2 gene in the Prader-Willi syndrome region is subject to genomic imprinting and positive selection. *Neurogenetics* 11: 153-161.

CHROMOSOMAL LOCATION

Genetic locus: CCDC33 (human) mapping to 15q24.1.

SOURCE

CCDC33 (F-3) is a mouse monoclonal antibody raised against amino acids 659-849 mapping near the C-terminus of CCDC33 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CCDC33 (F-3) is recommended for detection of CCDC33 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CCDC33 siRNA (h): sc-90062, CCDC33 shRNA Plasmid (h): sc-90062-SH and CCDC33 shRNA (h) Lentiviral Particles: sc-90062-V.

Molecular Weight of CCDC33: 114 kDa.


Positive Controls: CCDC33 (h): 293T Lysate: sc-114170.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.
- Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CCDC33 (F-3): sc-377074. Western blot analysis of CCDC33 expression in non-transfected (A) and human CCDC33 transfected (B) 293T whole cell lysates.



CCDC33 (F-3): sc-377074. Immunoperoxidase staining of formalin fixed, paraffin-embedded human prostate tissue showing cytoplasmic and membrane staining of glandular cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.