

JIP-2 (A-9): sc-377490

BACKGROUND

c-Jun NH₂-terminal kinases (JNKs) are distant members of the MAP kinase family. JNK1 is activated by dual phosphorylation at a Thr-Pro-Tyr motif in response to ultraviolet (UV) light, and it functions to phosphorylate c-Jun at amino terminal serine regulatory sites, Ser 63 and Ser 73, resulting in transcriptional activation. Two additional JNK family members have been identified as JNK2 and JNK3. JIP-1 (for JNK interacting protein-1) has been identified as a cytoplasmic inhibitor of JNK that retains JNK in the cytoplasm, thereby inhibiting JNK-regulated gene expression. Evidence suggests that JNK1 and JNK2 bind to JIP-1 with greater affinity than to ATF-2 and c-Jun, which are targets of the JNK signaling pathway. JIP-1 contains an amino terminal JNK binding domain and a carboxy terminal SH3 domain. ATF-2 and c-Jun also contain the JNK binding domain and are thought to compete with JIP-1 for JNK binding. Multiple splice variants of JIP-1, including JIP-1b, JIP-1c (also designated islet-brain 1 or IB-1), JIP-2a, JIP-2b and JIP-3, have been identified in brain.

REFERENCES

1. Pulverer, B.J., et al. 1991. Phosphorylation of c-Jun mediated by MAP kinases. *Nature* 353: 670-674.
2. Smeal, T., et al. 1992. Oncoprotein-mediated signalling cascade stimulates c-Jun activity by phosphorylation of serines 63 and 73. *Mol. Cell. Biol.* 12: 3507-3512.
3. Derijard, B., et al. 1994. JNK1: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* 76: 1025-1037.

CHROMOSOMAL LOCATION

Genetic locus: MAPK8IP2 (human) mapping to 22q13.33; Mapk8ip2 (mouse) mapping to 15 E3.

SOURCE

JIP-2 (A-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 51-89 near the N-terminus of JIP-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

JIP-2 (A-9) is available conjugated to agarose (sc-377490 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-377490 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-377490 PE), fluorescein (sc-377490 FITC), Alexa Fluor® 488 (sc-377490 AF488), Alexa Fluor® 546 (sc-377490 AF546), Alexa Fluor® 594 (sc-377490 AF594) or Alexa Fluor® 647 (sc-377490 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-377490 AF680) or Alexa Fluor® 790 (sc-377490 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-377490 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

JIP-2 (A-9) is recommended for detection of JIP-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

JIP-2 (A-9) is also recommended for detection of JIP-2 in additional species, including equine and canine.

Suitable for use as control antibody for JIP-2 siRNA (h): sc-40719, JIP-2 siRNA (m): sc-40720, JIP-2 shRNA Plasmid (h): sc-40719-SH, JIP-2 shRNA Plasmid (m): sc-40720-SH, JIP-2 shRNA (h) Lentiviral Particles: sc-40719-V and JIP-2 shRNA (m) Lentiviral Particles: sc-40720-V.


Molecular Weight of JIP-2: 88 kDa.

Positive Controls: JIP-2 (m): 293T Lysate: sc-121154, mouse brain extract: sc-2253 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



JIP-2 (A-9): sc-377490. Western blot analysis of JIP-2 expression in non-transfected: sc-117752 (A) and mouse JIP-2 transfected: sc-121154 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.