

p-Rsk-2 (C-6): sc-377501

BACKGROUND

The family of ribosomal S6 kinases (Rsk), designated Rsk-1 (or MAPKAP kinase-1), Rsk-2 and Rsk-3, are intracellular serine/threonine kinases that are important signaling intermediates in response to a broad range of ligand activated receptor tyrosine kinases. A unique feature common to the members of the Rsk family is that each possesses two non-identical complete kinase catalytic domains. An additional Rsk protein, Rsk-4, shows a high level of homology to the three previously isolated members of the human Rsk family. Rsk-4 is most abundantly expressed in brain and kidney and plays a role in normal neuronal development. The family of ribosomal S6 kinases includes p70 S6 kinase and p70 S6 kinase β , which are thought to have similar regulatory functions. MSK1 (also designated RLPK) is a novel Rsk-related protein, which, like the p90 Rsk family members, contains two non-identical complete kinase catalytic domains.

REFERENCES

- Alcorta, D.A., et al. 1989. Sequence and expression of chicken and mouse Rsk: homologs of *Xenopus laevis* ribosomal S6 kinase. *Mol. Cell. Biol.* 9: 3850-3859.
- Kozma, S.C., et al. 1990. Cloning of the mitogen-activated S6 kinase from rat liver reveals an enzyme of the second messenger subfamily. *Proc. Natl. Acad. Sci. USA* 87: 7365-7369.

CHROMOSOMAL LOCATION

Genetic locus: RPS6KA3 (human) mapping to Xp22.12; Rps6ka3 (mouse) mapping to X F4.

SOURCE

p-Rsk-2 (C-6) is a mouse monoclonal antibody epitope corresponding to a short amino acid sequence containing Thr 577 phosphorylated Rsk-2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

p-Rsk-2 (C-6) is available conjugated to agarose (sc-377501 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-377501 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-377501 PE), fluorescein (sc-377501 FITC), Alexa Fluor[®] 488 (sc-377501 AF488), Alexa Fluor[®] 546 (sc-377501 AF546), Alexa Fluor[®] 594 (sc-377501 AF594) or Alexa Fluor[®] 647 (sc-377501 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-377501 AF680) or Alexa Fluor[®] 790 (sc-377501 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-377501 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p-Rsk-2 (C-6) is recommended for detection of Thr 577 phosphorylated Rsk-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


p-Rsk-2 (C-6) is also recommended for detection of correspondingly phosphorylated Rsk-2 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Rsk-2 siRNA (h): sc-36441, Rsk-2 siRNA (m): sc-36442, Rsk-2 shRNA Plasmid (h): sc-36441-SH, Rsk-2 shRNA Plasmid (m): sc-36442-SH, Rsk-2 shRNA (h) Lentiviral Particles: sc-36441-V and Rsk-2 shRNA (m) Lentiviral Particles: sc-36442-V.


Molecular Weight of p-Rsk-2: 80 kDa.

Positive Controls: NIH/3T3 + UV cell lysate: sc-3804 or HeLa + PMA cell lysate: sc-2258.

DATA



Western blot analysis of Rsk-2 phosphorylation in untreated (A, D), PMA treated (B, E) and PMA and lambda protein phosphatase (sc-200312A) treated (C, F) HeLa whole cell lysates. Antibodies tested include p-Rsk-2 (C-6): sc-377501 (A, B, C) and Rsk-2 (C-19): sc-1430 (D, E, F).



p-Rsk-2 (C-6): sc-377501. Immunoperoxidase staining of formalin fixed, paraffin-embedded human colon tissue showing nuclear staining of glandular cells and endothelial cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human gall bladder tissue showing nuclear staining of glandular cells (B).

SELECT PRODUCT CITATIONS

- Lee, C.J., et al. 2018. Kaempferol targeting on the fibroblast growth factor receptor 3-ribosomal S6 kinase 2 signaling axis prevents the development of rheumatoid arthritis. *Cell Death Dis.* 9: 401.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.