# Egr-4 siRNA (m): sc-37830



The Power to Question

#### **BACKGROUND**

Egr-1, Egr-2, Egr-3 and Egr-4 are nuclear transcription factors belonging to the Egr C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family and containing three C<sub>2</sub>H<sub>2</sub>-type zinc fingers. As immediate early proteins, Egr transcription factors are rapidly induced by diverse extracellular stimuli. They are subject to tight differential control through diverse mechanisms at several levels of regulation: transcriptional; translational and posttranslational (including glycosylation, phosphorylation and redox) mechanisms; and protein-protein interaction. Egr-1 binds to the DNA sequence 5'-CGCCCCGC-3' (Egr-site), thereby activating transcription of target genes whose products are required for mitogenisis and differentiation. Egr-2 binds specific DNA sites located in the promoter region of HoxA4. Egr-2 defects cause congenital hypomyelination neuropathy (also designated Charcot-Marie-Tooth disease) and Dejerine-Sottas neuropathology (also designated hereditary motor and sensory neuropathy III). Egr-3 is involved in muscle spindle development and is expressed in T cells 20 minutes following activation. Egr-4 binds to the Egr consensus motif GCGTGGGCG, functions as a transcriptional repressor, and displays autoregulatory activities, downregulating its own gene promoter in a dose dependent manner.

## **REFERENCES**

- Beckmann, A. M. and Wilce, P. A. 1997. Egr transcription factors in the nervous system. Neurochem. Int. 31: 477-510.
- Zipfel, P.F., Decker, E.L., Holst, C. and Skerka C. 1997. The human zinc finger protein Egr-4 acts as autoregulatory transcriptional repressor. Biochim. Biophys. Acta 1354: 134-144.
- 3. SWISS-PROT/TrEMBL (P18146). World Wide Web URL: http://www.uniprot.org/uniprot/P18146

# CHROMOSOMAL LOCATION

Genetic locus: Egr4 (mouse) mapping to 6 C3.

# **PRODUCT**

Egr-4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu M$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Egr-4 shRNA Plasmid (m): sc-37830-SH and Egr-4 shRNA (m) Lentiviral Particles: sc-37830-V as alternate gene silencing products.

For independent verification of Egr-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-37830A, sc-37830B and sc-37830C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNAse-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

#### **APPLICATIONS**

Egr-4 siRNA (m) is recommended for the inhibition of Egr-4 expression in mouse cells.

## **SUPPORT REAGENTS**

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## **RT-PCR REAGENTS**

Semi-quantitative RT-PCR may be performed to monitor Egr-4 gene expression knockdown using RT-PCR Primer: Egr-4 (m)-PR: sc-37830-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com