

# GABP- $\beta$ 1/2 siRNA (h): sc-37903

## BACKGROUND

The transcription factor GA-binding protein (GABP) is composed of two subunits, the Ets-related GABP- $\alpha$  and a GABP- $\alpha$ -associated subunit, GABP- $\beta$ . GABP- $\alpha$  binds to a specific DNA sequence and GABP- $\beta$  exists as  $\beta$ 1 and  $\beta$ 2 splice variants that differ in their C-termini. In primary neuronal cultures, GABP- $\beta$  is expressed in both the cytoplasm and the nucleus, whereas GABP- $\alpha$  is expressed mainly in the nucleus. GABP is constitutively expressed as either a GABP- $\alpha\beta$  heterodimer or a GABP- $\alpha\beta$  heterotetramer, both of which can modify GABP-dependent transcription *in vitro* and *in vivo*. The GABP- $\alpha\beta$  tetrameric complex performs many different functions, such as stimulating transcription of the adenovirus E4 gene, differentially activating BRCA1 expression in human breast cell lines, potentiating Tat-mediated activation of long terminal repeat promoter transcription and viral replication in certain cell types, acting as a coordinator of mitochondrial and nuclear transcription for cytochrome oxidase in neurons and assisting in the regulation of rpl32 gene transcription.

## REFERENCES

1. Martin, M.E., et al. 1996. Redox regulation of GA-binding protein- $\alpha$  DNA binding activity. *J. Biol. Chem.* 271: 25617-25623.
2. Suzuki, F., et al. 1998. Functional interactions of transcription factor human GA-binding protein subunits. *J. Biol. Chem.* 273: 29302-29308.
3. Sawada, J., et al. 1999. Synergistic transcriptional activation by hGABP and select members of the activation transcription factor/cAMP response element-binding protein family. *J. Biol. Chem.* 274: 35475-35482.
4. Verhoef, K., et al. 1999. Evolution of the human immunodeficiency virus type 1 long terminal repeat promoter by conversion of an NF $\kappa$ B enhancer element into a GABP binding site. *J. Virol.* 73: 1331-1340.
5. Hoare, S., et al. 1999. Identification of a GABP  $\alpha/\beta$  binding site involved in the induction of oxytocin receptor gene expression in human breast cells, potentiation by c-Fos/c-Jun. *Endocrinology* 140: 2268-2279.
6. Chinenov, Y., et al. 2000. The  $\alpha$  and  $\beta$  subunits of the GA-binding protein form a stable heterodimer in solution. Revised model of heterotetrameric complex assembly. *J. Biol. Chem.* 275: 7749-7756.

## CHROMOSOMAL LOCATION

Genetic locus: GABPB1 (human) mapping to 15q21.2.

## PRODUCT

GABP- $\beta$ 1/2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GABP- $\beta$ 1/2 shRNA Plasmid (h): sc-37903-SH and GABP- $\beta$ 1/2 shRNA (h) Lentiviral Particles: sc-37903-V as alternate gene silencing products.

For independent verification of GABP- $\beta$ 1/2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-37903A, sc-37903B and sc-37903C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

GABP- $\beta$ 1/2 siRNA (h) is recommended for the inhibition of GABP- $\beta$ 1/2 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

GABP- $\beta$ 1/2 (E-7): sc-271571 is recommended as a control antibody for monitoring of GABP- $\beta$ 1/2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GABP- $\beta$ 1/2 gene expression knockdown using RT-PCR Primer: GABP- $\beta$ 1/2 (h)-PR: sc-37903-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.