

# Liprin $\beta$ 1 siRNA (m): sc-38032

## BACKGROUND

Liprins interact with members of the leukocyte common antigen-related (LAR) family of transmembrane protein tyrosine phosphatases, which are implicated in axon guidance and mammary gland development. Liprins are multivalent proteins that form complex structures and act as scaffolds for the recruitment and anchoring of LAR phosphatases. Based on sequence similarities and binding characteristics, liprins are subdivided into  $\alpha$ -type and  $\beta$ -type liprins. Both  $\alpha$ - and  $\beta$ -Liprins homodimerize via their N-terminal, coiled coil regions. Liprin  $\beta$ 1 interacts with S100A4, a calcium-binding protein related to tumor invasiveness and metastasis. The two proteins colocalize in the cytoplasm and at protrusion sites in the plasma membrane. The interaction of Liprin  $\beta$ 1 and S100A4 inhibits the phosphorylation of Liprin  $\beta$ 1 by PKC and protein kinase CK2 *in vitro*.

## REFERENCES

1. Serra-Pages, C., Medley, Q.G., Tang, M., Hart, A. and Streuli, M. 1998. Liprins, a family of LAR transmembrane protein-tyrosine phosphatase-interacting proteins. *J. Biol. Chem.* 273: 15611-15620.
2. Kriajevska, M., Fischer-Larsen, M., Moertz, E., Vorm, O., Tulchinsky, E., Grigorian, M., Ambartsumian, N. and Lukanidin, E. 2002. Liprin  $\beta$ 1, a member of the family of LAR transmembrane tyrosine phosphatase-interacting proteins, is a new target for the metastasis-associated protein S100A4 (Mts1). *J. Biol. Chem.* 277: 5229-5235.
3. Katoh, M. 2003. Identification and characterization of human PPFIA4 gene *in silico*. *Int. J. Mol. Med.* 12: 1009-1014.

## CHROMOSOMAL LOCATION

Genetic locus: Ppfbp1 (mouse) mapping to 6 G3.

## PRODUCT

Liprin  $\beta$ 1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Liprin  $\beta$ 1 shRNA Plasmid (m): sc-38032-SH and Liprin  $\beta$ 1 shRNA (m) Lentiviral Particles: sc-38032-V as alternate gene silencing products.

For independent verification of Liprin  $\beta$ 1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38032A, sc-38032B and sc-38032C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Liprin  $\beta$ 1 siRNA (m) is recommended for the inhibition of Liprin  $\beta$ 1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

Liprin  $\beta$ 1 (F-2): sc-514575 is recommended as a control antibody for monitoring of Liprin  $\beta$ 1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Liprin  $\beta$ 1 gene expression knockdown using RT-PCR Primer: Liprin  $\beta$ 1 (m)-PR: sc-38032-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.