

SLBP siRNA (h): sc-38321

BACKGROUND

Replication-dependent histone mRNAs lack polyadenylated tails and instead end in a conserved stem-loop. The stem-loop binding protein (SLBP) binds the 3' end of histone mRNA and contains a 73 amino acid RNA-binding domain. SLBP mediates the interaction of the histone pre-mRNA with U7 SnRNP to facilitate 3' end processing. SLBP is required for the translation of stem-loop mRNAs. SLBP forms a stable complex with U7 snRNP in the nucleus as well as the cytoplasm. hZFP100 is a zinc finger protein that interacts with the SLBP/RNA complex but not with free SLBP. During the cell cycle, SLBP increases in the late G₁ and decreases in the S/G₂ border. The regulation of SLBP occurs at the level of translation. Specifically, two phosphorylation events on Threonine 99 and Threonine 104 trigger the degradation of SLBP in late S-phase cells.

REFERENCES

1. Wang, Z.F., et al. 1996. The protein that binds the 3' end of histone mRNA: a novel RNA-binding protein required for histone pre-mRNA processing. *Genes Dev.* 10: 3028-3040.
2. Martin, F., et al. 1997. The gene for histone RNA hairpin binding protein is located on human chromosome 4 and encodes a novel type of RNA binding protein. *EMBO J.* 16: 769-778.
3. Dominski, Z., et al. 1999. Stem-loop binding protein facilitates 3' end formation by stabilizing U7 snRNP binding to histone pre-mRNA. *Mol. Cell. Biol.* 19: 3561-3570.
4. Whitfield, M.L., et al. 2000. Stem-loop binding protein, the protein that binds the 3' end of histone mRNA, is cell cycle regulated by both translational and posttranslational mechanisms. *Mol. Cell. Biol.* 20: 4188-4198.
5. Ling, J., et al. 2002. The histone 3' terminal stem-loop-binding protein enhances translation through a functional and physical interaction with eukaryotic initiation factor 4G (eIF4G) and eIF3. *Mol. Cell. Biol.* 22: 7853-7867.
6. Dominski, Z., et al. 2002. A novel zinc finger protein is associated with U7 snRNP and interacts with the stem-loop binding protein in the histone pre-mRNA to stimulate 3' end processing. *Genes Dev.* 16: 58-71.

CHROMOSOMAL LOCATION

Genetic locus: SLBP (human) mapping to 4p16.3.

PRODUCT

SLBP siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SLBP shRNA Plasmid (h): sc-38321-SH and SLBP shRNA (h) Lentiviral Particles: sc-38321-V as alternate gene silencing products.

For independent verification of SLBP (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38321A, sc-38321B and sc-38321C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SLBP siRNA (h) is recommended for the inhibition of SLBP expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

SLBP (H-3): sc-376310 is recommended as a control antibody for monitoring of SLBP gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SLBP gene expression knockdown using RT-PCR Primer: SLBP (h)-PR: sc-38321-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.