

Sm D1 siRNA (h): sc-38325

BACKGROUND

U1, U2, U4, U5, U6 and U7 are small nuclear ribonucleoproteins (snRNPs) that comprise the spliceosome in eukaryotes. Each snRNP contains common Sm proteins B/B', D1, D2, D3, E, F and G. The Sm proteins pair up as D1-D2, B/B'-D3 and E-F-G to form RNA-free hetero-oligomers in the cytoplasm. Sm proteins aid in the cytoplasmic construction of the snRNPs by binding to a conserved Sm site on snRNA and forming a stable snRNP core complex. Sm D1, D2 and D3 are present in U1, U2, U4/5 and U5 but not U7 snRNPs in human and mouse cells. U7 snRNPs contain Lsm10, an Sm D1-like protein. Autoantibodies produced in patients suffering from systemic lupus erythematosus react predominantly with Sm B/B', D1 and D3. The major linear epitope of these autoantibodies includes the C-terminal RG dipeptide repeats found in Sm D1 and D3.

REFERENCES

1. Branlant, C., Krol, A., Ebel, J.P., Lazar, E., Haendler, B. and Jacob, M. 1982. U2 RNA shares a structural domain with U1, U4, and U5 RNAs. *EMBO J.* 1: 1259-1265.
2. Lehmeier, T., Foulaki, K. and Luhrmann, R. 1990. Evidence for three distinct D proteins, which react differentially with anti-Sm autoantibodies, in the cores of the major snRNPs U1, U2, U4/U6 and U5. *Nucleic Acids Res.* 18: 6475-6484.
3. Raker, V.A., Plessel, G. and Luhrmann, R. 1996. The snRNP core assembly pathway: identification of stable core protein heteromeric complexes and an snRNP subcore particle *in vitro*. *EMBO J.* 15: 2256-2269.
4. Brahms, H., Raymackers, J., Union, A., Keyser, F., Meheus, L. and Luhrmann, R. 2000. The C-terminal RG dipeptide repeats of the spliceosomal Sm proteins D1 and D3 contain symmetrical dimethylarginines, which form a major B-cell epitope for anti-Sm autoantibodies. *J. Biol. Chem.* 275: 17122-17129.
5. Pillai, R.S., Will, C.L., Luhrmann, R., Schumperli, D. and Muller, B. 2001. Purified U7 snRNPs lack the Sm proteins D1 and D2 but contain Lsm10, a new 14 kDa Sm D1-like protein. *EMBO J.* 20: 5470-5479.
6. LocusLink Report (LocusID: 6632). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: SNRPD1 (human) mapping to 18q11.2.

PRODUCT

Sm D1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Sm D1 shRNA Plasmid (h): sc-38325-SH and Sm D1 shRNA (h) Lentiviral Particles: sc-38325-V as alternate gene silencing products.

For independent verification of Sm D1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38325A, sc-38325B and sc-38325C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Sm D1 siRNA (h) is recommended for the inhibition of Sm D1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Sm D1 (A-9): sc-166650 is recommended as a control antibody for monitoring of Sm D1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Sm D1 gene expression knockdown using RT-PCR Primer: Sm D1 (h)-PR: sc-38325-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.