

Smad3 siRNA (h): sc-38376

BACKGROUND

Smad proteins, the mammalian homologs of the *Drosophila* mothers against dpp (Mad) have been implicated as downstream effectors of TGF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1), Smad5 and mammalian Smad8 (also designated Smad9 or MADH6) are effectors of BMP2 and BMP4 function while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad family members.

CHROMOSOMAL LOCATION

Genetic locus: SMAD3 (human) mapping to 15q22.33.

PRODUCT

Smad3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Smad3 shRNA Plasmid (h): sc-38376-SH and Smad3 shRNA (h) Lentiviral Particles: sc-38376-V as alternate gene silencing products.

For independent verification of Smad3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38376A, sc-38376B and sc-38376C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Smad3 siRNA (h) is recommended for the inhibition of Smad3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Smad2/3 (C-8): sc-133098 is recommended as a control antibody for monitoring of Smad3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Smad3 gene expression knockdown using RT-PCR Primer: Smad3 (h)-PR: sc-38376-PR (20 μ l, 495 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Nold, M.F., et al. 2010. IL-37 is a fundamental inhibitor of innate immunity. *Nat. Immunol.* 11: 1014-1022.
- Chowdhury, S., et al. 2011. Identification of a novel TGF β /PKA signaling transduceome in mediating control of cell survival and metastasis in colon cancer. *PLoS ONE* 6: e19335.
- Fang, W.B., et al. 2012. CCL2/CCR2 chemokine signaling coordinates survival and motility of breast cancer cells through Smad3 protein- and p42/44 mitogen-activated protein kinase (MAPK)-dependent mechanisms. *J. Biol. Chem.* 287: 36593-36608.
- Qin, H., et al. 2013. Ossifying fibroma tumor stem cells are maintained by epigenetic regulation of a TSP1/TGF- β /Smad3 autocrine loop. *Cell Stem Cell* 13: 577-589.
- Lin, L., et al. 2014. Transcriptional regulation of Stat3 by SPTBN1 and Smad3 in HCC through cAMP-response element-binding proteins ATF3 and CREB2. *Carcinogenesis* 35: 2393-2403.
- Yu, Y. and Ran, Q. 2015. Nuclear Smad2 restrains proliferation of glioblastoma. *Cell. Physiol. Biochem.* 35: 1756-1763.
- Chen, J., et al. 2016. TGF- β / β 2-spectrin/CTCF-regulated tumor suppression in human stem cell disorder Beckwith-Wiedemann syndrome. *J. Clin. Invest.* 126: 527-542.
- Chen, J., et al. 2016. Vitamin D deficiency promotes liver tumor growth in transforming growth factor- β /Smad3-deficient mice through Wnt and Toll-like receptor 7 pathway modulation. *Sci. Rep.* 6: 30217.
- Zhang, Z., et al. 2018. Correction: opposing effects of PI3K/Akt and Smad-dependent signaling pathways in NAG-1-induced glioblastoma cell apoptosis. *PLoS ONE* 13: e0205391.
- Rahn, S., et al. 2018. Diabetes as risk factor for pancreatic cancer: hyperglycemia promotes epithelial-mesenchymal-transition and stem cell properties in pancreatic ductal epithelial cells. *Cancer Lett.* 415: 129-150.
- Han, R., et al. 2019. C3a and suPAR drive versican V1 expression in tubular cells of focal segmental glomerulosclerosis. *JCI Insight* 4: e122912.

RESEARCH USE

For research use only, not for use in diagnostic procedures.