



GBX2 siRNA (m): sc-38666

BACKGROUND

The isthmic organizer signals at the mid/hindbrain boundary (MHB) regulate the development and differentiation of the vertebrate caudal midbrain and the anterior hindbrain. The MHB forms at the boundary of expression between homeobox genes *Gbx2* and *Otx2*. *Gbx2* and *Otx2* play distinct, essential roles in MHB positioning and development. During development, the *GBX2* gene is expressed in the anterior hindbrain. Specifically, *Gbx2* negatively regulates *Otx2* expression along the anterior-posterior axis; *Gbx2*(-) mutants demonstrate an expanded *Otx2* domain. During development, the *GBX2* gene is expressed in the anterior hindbrain. *Gbx2* is expressed in the adult brain, spleen and female genital tract. The *GBX2* gene is over-expressed in human prostate cancer cell lines (TSU-prl, PC3, DU145 and LNCaP). Furthermore, downregulation of *Gbx2* expression restricts tumorigenicity in human prostate cancer cell lines, which suggests that *Gbx2* expression may be required for growth of malignant prostate cells.

REFERENCES

1. Bouillet, P., et al. 1995. Sequence and expression pattern of the *Stra7* (*Gbx-2*) homeobox-containing gene induced by retinoic acid in P19 embryonal carcinoma cells. *Dev. Dyn.* 204: 372-382.
2. Gao, A.C. et al. 1996. Expression of homeobox gene-*GBX2* in human prostatic cancer cells. *Prostate* 29: 395-398.
3. Gao, A.C., et al. 1998. Downregulation of homeobox gene *GBX2* expression inhibits human prostate cancer clonogenic ability and tumorigenicity. *Cancer Res.* 58: 1391-1394.
4. Millet, S., et al. 1999. A role for *Gbx2* in repression of *Otx2* and positioning the mid/hindbrain organizer. *Nature* 401: 161-164.
5. Gao, A.C., et al. 2000. Enhanced *GBX2* expression stimulates growth of human prostate cancer cells via transcriptional upregulation of the Interleukin 6 gene. *Clin. Cancer Res.* 6: 493-497.
6. Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 601135. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Tour, E., et al. 2002. *Otx2* can activate the isthmic organizer genetic network in the *Xenopus* embryo. *Mech. Dev.* 110: 3-13.

CHROMOSOMAL LOCATION

Genetic locus: *Gbx2* (mouse) mapping to 1 D.

PRODUCT

GBX2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see *GBX2* shRNA Plasmid (m): sc-38666-SH and *GBX2* shRNA (m) Lentiviral Particles: sc-38666-V as alternate gene silencing products.

For independent verification of *GBX2* (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38666A, sc-38666B and sc-38666C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GBX2 siRNA (m) is recommended for the inhibition of *GBX2* expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor *GBX2* gene expression knockdown using RT-PCR Primer: *GBX2* (m)-PR: sc-38666-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.