

HoxA7 siRNA (h): sc-38680

BACKGROUND

HOX genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. The homeobox gene HoxA1 is transcriptionally regulated by retinoic acid (RA) and encodes a transcription factor, which has been shown to play important roles in cell differentiation and embryogenesis. HoxA1 is also expressed in cancers, such as mammary tumors, though it is not expressed in normal gland or in precancerous mammary tissues. At embryonic stages, HoxA2 is expressed in the mesenchyme and epithelial cells of palate, however its expression is restricted to the tips of the growing palatal shelves. HoxA2 protein is predominantly expressed in the nuclei of cells in the ventral mantle region of the developing embryo. In the developing and adult mouse spinal cord, HoxA2 protein may contribute to dorsal-ventral patterning and/or to the specification of neuronal phenotype. HoxA7 functions as a potent transcriptional repressor and its action as such requires several domains, including both activator and repressor regions. HoxA7 is expressed in the fetal liver, lung, skeletal muscle, kidney, pancreas and placenta.

REFERENCES

1. Schnabel, C.A. et al. 1996. Repression by HoxA7 is mediated by the homeodomain and the modulatory action of its N-terminal-arm residues. *Mol. Cell. Biol.* 16: 2678-2688.
2. Srebrow, A., et al. 1998. Expression of HoxA1 and HoxB7 is regulated by extracellular matrix-dependent signals in mammary epithelial cells. *J. Cell Biol.* 69: 377-391.
3. Hao, Z., et al. 1999. Differential expression of HoxA2 protein along the dorsal-ventral axis of the developing and adult mouse spinal cord. *Dev. Dyn.* 216: 201-217.
4. Kim, M.H., et al. 2000. Sequence analysis and tissue specific expression of human HoxA7. *Mol. Biotechnol.* 14: 19-24.
5. Nazarali, A., et al. 2000. Temporal and spatial expression of HoxA2 during murine palatogenesis. *Cell. Mol. Neurobiol.* 20: 269-290.
6. Shen, J., et al. 2000. Molecular cloning and analysis of a group of genes differentially expressed in cells which overexpress the HoxA1 homeobox gene. *Exp. Cell Res.* 259: 274-283.
7. Goodman, F.R., et al. 2001. Human HOX gene mutations. *Clin. Genet.* 59: 1-11.

CHROMOSOMAL LOCATION

Genetic locus: HOXA7 (human) mapping to 7p15.2.

PRODUCT

HoxA7 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HoxA7 shRNA Plasmid (h): sc-38680-SH and HoxA7 shRNA (h) Lentiviral Particles: sc-38680-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HoxA7 siRNA (h) is recommended for the inhibition of HoxA7 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

HoxA7 (743C1a): sc-81290 is recommended as a control antibody for monitoring of HoxA7 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HoxA7 gene expression knockdown using RT-PCR Primer: HoxA7 (h)-PR: sc-38680-PR (20 μ l, 420 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.