

TRβ1 siRNA (h): sc-38890

BACKGROUND

Thyroid hormone nuclear receptors (TRs) are ligand-dependent transcription factors which regulate growth, differentiation and development, and represent members of the steroid/retinoic acid superfamily. The two genes encoding TRs identified to date, TRα and TRβ, have been mapped to human chromosomes 17 and 3, respectively. TRs bind to thyroid hormone response elements (TREs) with half-site binding motifs in the orientation of palindromes, direct repeats or inverted palindromes. The affinities of binding are both variable and influenced differentially by 3,5,3'-triiodo-L-thyronine (T3). Transcriptional regulation by TRs is also modulated by heterodimerization with TR nuclear accessory proteins, the most extensively characterized of which are the retinoid X receptors (RXRα, RXRβ and RXRγ). The TRβ isoform TRβ1 forms a complex with the PI 3-kinase p85α subunit and plays an important role in the T3-induced activation of Akt in pancreatic β cells.

REFERENCES

- Näär, A., et al. 1991. The orientation and spacing of core DNA-binding motifs dictate selective transcriptional responses to three nuclear receptors. *Cell* 65: 1267-1271.
- Lazar, M.A. 1993. Thyroid hormone receptors: multiple forms, multiple possibilities. *Endocrinol. Rev.* 14: 184-193.
- Meier, C.A., et al. 1993. Interaction of human TRβ1 and its mutants with DNA and RXRβ. T3 response element-dependent dominant negative potency. *J. Clin. Invest.* 92: 1986-1993.
- Zhang, X.K., et al. 1993. Hetero- and homodimeric receptors in thyroid hormone and vitamin A action. *Receptor* 3: 183-191.
- Bhat, M.K., et al. 1994. Phosphorylation enhances the target gene sequence-dependent dimerization of thyroid hormone receptor with retinoid X receptor. *Proc. Natl. Acad. Sci. USA* 91: 7927-7931.
- Sugawara, A., et al. 1994. Phosphorylation selectively increases triiodo-thyronine receptor homodimer binding to DNA. *J. Biol. Chem.* 269: 433-437.

CHROMOSOMAL LOCATION

Genetic locus: THRB (human) mapping to 3p24.2.

PRODUCT

TRβ1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRβ1 shRNA Plasmid (h): sc-38890-SH and TRβ1 shRNA (h) Lentiviral Particles: sc-38890-V as alternate gene silencing products.

For independent verification of TRβ1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38890A, sc-38890B and sc-38890C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRβ1 siRNA (h) is recommended for the inhibition of TRβ1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TRβ1 (J51): sc-737 is recommended as a control antibody for monitoring of TRβ1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRβ1 gene expression knockdown using RT-PCR Primer: TRβ1 (h)-PR: sc-38890-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Flamini, M.I., et al. 2017. Thyroid hormone controls breast cancer cell movement via integrin α_v/β₃/SRC/FAK/PI3-kinases. *Horm. Cancer* 8: 16-27.

RESEARCH USE

For research use only, not for use in diagnostic procedures.