

AK4 siRNA (m): sc-38909

BACKGROUND

Adenylate kinases 1-5 (designated AK1-5) are a set of enzymes that regulate the phosphorylation state of intracellular adenine nucleotides, which are the principle high-energy phosphoryl-carrying molecules in living cells. AKs influence metabolic signals, which include gene expression, ion channel activity and protein kinase-mediated signaling, by catalyzing phosphoryl transfer between adenine nucleotides (AMP, ADP, ATP). Inherited mutations leading to AK deficiencies in erythrocytes have been implicated in hemolytic anemia. Rat AK4 mRNA is expressed as a 223 amino acid protein in the central nervous system from the middle stage of embryogenesis to adulthood.

REFERENCES

1. Dzeja, P.P., Zeleznikar, R.J. and Goldberg, N.D. 1998. Adenylate kinase: kinetic behavior in intact cells indicates it is integral to multiple cellular processes. *Mol. Cell. Biochem.* 184: 169-182.
2. Yoneda, T., Sato, M., Maeda, M. and Takagi, H. 1998. Identification of a novel adenylate kinase system in the brain: cloning of the fourth adenylate kinase. *Brain Res. Mol. Brain Res.* 62: 187-195.
3. Online Mendelian Inheritance in Man, OMIM[™]. 1999. Johns Hopkins University, Baltimore, MD. MIM Number: 103000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Carrasco, A.J., Dzeja, P.P., Alekseev, A.E., Pucar, D., Zingman, L.V., Abraham, M.R., Hodgson, D., Bienengraeber, M., Puceat, M., Janssen, E., Wieringa, B. and Terzic, A. 2001. Adenylate kinase phosphotransfer communicates cellular energetic signals to ATP-sensitive potassium channels. *Proc. Natl. Acad. Sci. USA* 98: 7623-7628.
5. LocusLink Report (LocusID: 205). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: Ak4 (mouse) mapping to 4 C6.

PRODUCT

AK4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AK4 shRNA Plasmid (m): sc-38909-SH and AK4 shRNA (m) Lentiviral Particles: sc-38909-V as alternate gene silencing products.

For independent verification of AK4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38909A, sc-38909B and sc-38909C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AK4 siRNA (m) is recommended for the inhibition of AK4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

AK4 (A-9): sc-271161 is recommended as a control antibody for monitoring of AK4 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AK4 gene expression knockdown using RT-PCR Primer: AK4 (m)-PR: sc-38909-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.