

Akt3 siRNA (h): sc-38911

BACKGROUND

The serine/threonine kinase Akt family contains several members, including Akt1 (also designated PKB or RacPK), Akt2 (also designated PKB β or RacPK- β) and Akt3 (also designated PKB γ or thymoma viral proto-oncogene 3), which exhibit sequence homology with the protein kinase A and C families and are encoded by the *c-Akt* proto-oncogene. All members of the Akt family have a pleckstrin homology domain. Akt3 is phosphorylated on a serine residue in response to Insulin. However, the activation of Akt3 by Insulin is inhibited by prior activation of protein kinase C via a mechanism that does not require the presence of the PH domain. Akt3 is expressed in 3T3-L1 fibroblasts, adipocytes and skeletal muscle and may be involved in various biological processes, including adipocyte and muscle differentiation, glycogen synthesis, glucose uptake, apoptosis and cellular proliferation.

CHROMOSOMAL LOCATION

Genetic locus: AKT3 (human) mapping to 1q43.

PRODUCT

Akt3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Akt3 shRNA Plasmid (h): sc-38911-SH and Akt3 shRNA (h) Lentiviral Particles: sc-38911-V as alternate gene silencing products.

For independent verification of Akt3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38911A, sc-38911B and sc-38911C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Akt3 siRNA (h) is recommended for the inhibition of Akt3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Akt3 (EE-M14): sc-134254 is recommended as a control antibody for monitoring of Akt3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Akt3 gene expression knockdown using RT-PCR Primer: Akt3 (h)-PR: sc-38911-PR (20 μ l, 461 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

SELECT PRODUCT CITATIONS

- Toschi, A., et al. 2008. Differential dependence of hypoxia-inducible factors 1 α and 2 α on mTORC1 and mTORC2. *J. Biol. Chem.* 283: 34495-34499.
- Moro, L., et al. 2009. Mitochondrial DNA depletion in prostate epithelial cells promotes anoikis resistance and invasion through activation of PI3K/Akt2. *Cell Death Differ.* 16: 571-583.
- Moriya, C., et al. 2011. Expression of matrix metalloproteinase-13 is controlled by IL-13 via PI3K/Akt3 and PKC- δ in normal human dermal fibroblasts. *J. Invest. Dermatol.* 131: 655-661.
- Schultz, C.R., et al. 2012. Inhibition of HSP 27 alone or in combination with pAkt inhibition as therapeutic approaches to target SPARC-induced glioma cell survival. *Mol. Cancer* 11: 20.
- Lehman, H.L., et al. 2012. Regulation of inflammatory breast cancer cell invasion through Akt1/PKB α phosphorylation of RhoC GTPase. *Mol. Cancer Res.* 10: 1306-1318.
- Corum, D.G., et al. 2014. Akt3 controls mitochondrial biogenesis and autophagy via regulation of the major nuclear export protein CRM-1. *FASEB J.* 28: 395-407.
- Wang, J., et al. 2015. MicroRNA-29b prevents liver fibrosis by attenuating hepatic stellate cell activation and inducing apoptosis through targeting PI3K/Akt pathway. *Oncotarget* 6: 7325-7338.
- Rao, G., et al. 2017. Inhibition of Akt1 signaling promotes invasion and metastasis of non-small cell lung cancer cells with K-Ras or EGFR mutations. *Sci. Rep.* 7: 7066.
- Ding, L., et al. 2017. Akt3 kinase suppresses pinocytosis of low-density lipoprotein by macrophages via a novel WNK/SGK1/Cdc42 protein pathway. *J. Biol. Chem.* 292: 9283-9293.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.