

# AMPK $\beta$ 1 siRNA (m): sc-38926

## BACKGROUND

AMPK (for 5'-AMP-activated protein kinase) is a heterotrimeric complex comprising a catalytic  $\alpha$  subunit and regulatory  $\beta$  and  $\gamma$  subunits. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. AMPK is activated by high AMP and low ATP through a mechanism involving allosteric regulation, promotion of phosphorylation by an upstream protein kinase known as AMPK kinase and inhibition of dephosphorylation. Activated AMPK can phosphorylate and regulate *in vivo* hydroxymethylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively. The human AMPK $\alpha$ 1 and AMPK $\alpha$ 2 genes encode 548 amino acid and 552 amino acid proteins, respectively. Human AMPK $\beta$ 1 encodes a 271 amino acid protein and human AMPK $\beta$ 2 encodes a 272 amino acid protein. The human AMPK $\gamma$ 1 gene encodes a 331 amino acid protein. Human AMPK $\gamma$ 2 and AMPK $\gamma$ 3, which are 569 and 492 amino acid proteins, respectively, contain unique N-terminal domains and may participate directly in the binding of AMP within the AMPK complex.

## REFERENCES

1. Stapleton, D., et al. 1996. Mammalian AMP-activated protein kinase subfamily. *J. Biol. Chem.* 271: 611-614.
2. Stapleton, D., et al. 1997. AMP-activated protein kinase isoenzyme family: subunit structure and chromosomal location. *FEBS Lett.* 409: 452-456.
3. Hardie, D.G., et al. 1997. The AMP-activated protein kinase-fuel gauge of the mammalian cell? *Eur. J. Biochem.* 246: 259-273.
4. Thornton, C., et al. 1998. Identification of a novel AMP-activated protein kinase  $\beta$  subunit isoform that is highly expressed in skeletal muscle. *J. Biol. Chem.* 273: 12443-12450.
5. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 602739. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: Prkab1 (mouse) mapping to 5 F.

## PRODUCT

AMPK $\beta$ 1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AMPK $\beta$ 1 shRNA Plasmid (m): sc-38926-SH and AMPK $\beta$ 1 shRNA (m) Lentiviral Particles: sc-38926-V as alternate gene silencing products.

For independent verification of AMPK $\beta$ 1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38926A, sc-38926B and sc-38926C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

AMPK $\beta$ 1 siRNA (m) is recommended for the inhibition of AMPK $\beta$ 1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

AMPK $\beta$ 1 (Z14): sc-100357 is recommended as a control antibody for monitoring of AMPK $\beta$ 1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AMPK $\beta$ 1 gene expression knockdown using RT-PCR Primer: AMPK $\beta$ 1 (m)-PR: sc-38926-PR (20  $\mu$ l, 592 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.