

AMPK γ 1 siRNA (h): sc-38929

BACKGROUND

AMPK (for 5'-AMP-activated protein kinase) is a heterotrimeric complex comprising a catalytic α subunit and regulatory β and γ subunits. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. AMPK is activated by high AMP and low ATP through a mechanism involving allosteric regulation, promotion of phosphorylation by an upstream protein kinase known as AMPK kinase and inhibition of dephosphorylation. Activated AMPK can phosphorylate and regulate *in vivo* hydroxymethylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively. The human AMPK α 1 and AMPK α 2 genes encode 548 amino acid and 552 amino acid proteins, respectively. Human AMPK β 1 encodes a 271 amino acid protein and human AMPK β 2 encodes a 272 amino acid protein. The human AMPK γ 1 gene encodes a 331 amino acid protein. Human AMPK γ 2 and AMPK γ 3, which are 569 and 492 amino acid proteins, respectively, contain unique N-terminal domains and may participate directly in the binding of AMP within the AMPK complex.

REFERENCES

1. Stapleton, D., et al. 1996. Mammalian AMP-activated protein kinase subfamily. *J. Biol. Chem.* 271: 611-614.
2. Stapleton, D., et al. 1997. AMP-activated protein kinase isoenzyme family: subunit structure and chromosomal location. *FEBS Lett.* 409: 452-456.
3. Hardie, D.G., et al. 1997. The AMP-activated protein kinase-fuel gauge of the mammalian cell? *Eur. J. Biochem.* 246: 259-273.
4. Thornton, C., et al. 1998. Identification of a novel AMP-activated protein kinase β subunit isoform that is highly expressed in skeletal muscle. *J. Biol. Chem.* 273: 12443-12450.
5. Online Mendelian Inheritance in Man, OMIM[™]. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 602739. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Cheung, P.C., et al. 2000. Characterization of AMP-activated protein kinase γ subunit isoforms and their role in AMP binding. *Biochem. J.* 346: 659-669.

CHROMOSOMAL LOCATION

Genetic locus: PRKAG1 (human) mapping to 12q13.12.

PRODUCT

AMPK γ 1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AMPK γ 1 shRNA Plasmid (h): sc-38929-SH and AMPK γ 1 shRNA (h) Lentiviral Particles: sc-38929-V as alternate gene silencing products.

For independent verification of AMPK γ 1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38929A, sc-38929B and sc-38929C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AMPK γ 1 siRNA (h) is recommended for the inhibition of AMPK γ 1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AMPK γ 1 gene expression knockdown using RT-PCR Primer: AMPK γ 1 (h)-PR: sc-38929-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Bakula, D., et al. 2017. WIPI3 and WIPI4 β -propellers are scaffolds for LKB1-AMPK-TSC signalling circuits in the control of autophagy. *Nat. Commun.* 8: 15637.
2. Rosenthal, C.K. 2017. EGFR probes matrix stiffness. *Nat. Cell Biol.* 19: 600.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.